

## SIAE (S-19): sc-244135

### BACKGROUND

SIAE (sialic acid acetyltransferase) is a 523 amino acid secreted protein that is widely expressed with high expression in the testis, prostate and colon. Sialic acids are acidic 9-carbon sugars typically found at the nonreducing end of sugar chains. They are frequently modified by 9-O-acetylation, and this modification is removed by sialic acid acetyltransferases. The SIAE protein catalyzes the removal of O-acetyl ester groups from position 9 of the parent sialic acid, N-acetylneuraminic acid. Defects in SIAE are a cause of autoimmune disease type 6 (AIS6). Individuals manifesting susceptibility to AIS6 can suffer from juvenile idiopathic arthritis, rheumatoid arthritis, multiple sclerosis, Sjogren syndrome, systemic lupus erythematosus, type 1 diabetes, ulcerative colitis and Crohn disease. Existing as two alternatively spliced isoforms, the SIAE gene is conserved in chimpanzee, canine, bovine, mouse and zebrafish, and maps to human chromosome 11q24.2.

### REFERENCES

1. Takematsu, H., Diaz, S., Stoddart, A., Zhang, Y. and Varki, A. 1999. Lysosomal and cytosolic sialic acid 9-O-acetyltransferase activities can be encoded by one gene via differential usage of a signal peptide-encoding exon at the N terminus. *J. Biol. Chem.* 274: 25623-25631.
2. Zhu, H., Chan, H.C., Zhou, Z., Li, J., Zhu, H., Yin, L., Xu, M., Cheng, L. and Sha, J. 2004. A gene encoding sialic-acid-specific 9-O-acetyltransferase found in human adult testis. *J. Biomed. Biotechnol.* 2004: 130-136.
3. Taylor, T.D., Noguchi, H., Totoki, Y., Toyoda, A., Kuroki, Y., Dewar, K., Lloyd, C., Itoh, T., Takeda, T., Kim, D.W., She, X., Barlow, K.F., Bloom, T., Bruford, E., Chang, J.L., Cuomo, C.A., Eichler, E., FitzGerald, M.G., Jaffe, D.B., LaButti, K., Nicol, R., Park, H.S., et al. 2006. Human chromosome 11 DNA sequence and analysis including novel gene identification. *Nature* 440: 497-500.
4. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610079. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Cariappa, A., Takematsu, H., Liu, H., Diaz, S., Haider, K., Boboila, C., Kalloo, G., Connole, M., Shi, H.N., Varki, N., Varki, A. and Pillai, S. 2009. B cell antigen receptor signal strength and peripheral B cell development are regulated by a 9-O-acetyl sialic acid esterase. *J. Exp. Med.* 206: 125-138.
6. Suroli, I., Pirnie, S.P., Chellappa, V., Taylor, K.N., Cariappa, A., Moya, J., Liu, H., Bell, D.W., Driscoll, D.R., Diederichs, S., Haider, K., Netravali, I., Le, S., Elia, R., Dow, E., Lee, A., Freudenberg, J., De Jager, P.L., Chretien, Y., Varki, A., Macdonald, M.E., et al. 2010. Functionally defective germline variants of sialic acid acetyltransferase in autoimmunity. *Nature* 466: 243-247.

### CHROMOSOMAL LOCATION

Genetic locus: SIAE (human) mapping to 11q24.2.

### SOURCE

SIAE (S-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of SIAE of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244135 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

SIAE (S-19) is recommended for detection of SIAE of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SIAE siRNA (h): sc-96456, SIAE shRNA Plasmid (h): sc-96456-SH and SIAE shRNA (h) Lentiviral Particles: sc-96456-V.

Molecular Weight of SIAE isoforms 1/2: 58/55 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.