

SLC25A42 (T-16): sc-244163

BACKGROUND

SLC25A42 (solute carrier family 25, member 42) is a 318 amino acid multi-pass membrane protein that belongs to the mitochondrial carrier family. SLC25A42 contains three Solcar repeats. Expression of SLC25A42 was found in all rat tissues examined, with highest expression in adipose, followed by hypothalamus and brain coronal sections containing corpus callosum, fornix, thalamus, hypothalamus, optic chiasm, pons, midbrain and cerebellum. The main physiological role of SLC25A42 is to import CoA into mitochondria in exchange for intramitochondrial (deoxy)adenine nucleotides and adenosine 3',5'-diphosphate. SLC25A42 exhibits a high transport affinity for CoA, de-phospho-CoA, ADP and adenosine 3',5'-diphosphate. The SLC25A42 gene contains seven exons, is conserved in chimpanzee, canine, bovine, mouse, rat, chicken, zebrafish, fruit fly, mosquito, *C. elegans*, *S. pombe*, *S. cerevisiae*, *K. lactis*, *E. gossypii*, *M. grisea*, *N. crassa*, *A. thaliana* and rice, and maps to human chromosome 19p13.11. The mouse gene maps to chromosome 8.

REFERENCES

- Grimwood, J., Gordon, L.A., Olsen, A., Terry, A., Schmutz, J., Lamerdin, J., et al. 2004. The DNA sequence and biology of human chromosome 19. *Nature* 428: 529-535.
- Haitina, T., Lindblom, J., Renström, T. and Fredriksson, R. 2006. Fourteen novel human members of mitochondrial solute carrier family 25 (SLC25) widely expressed in the central nervous system. *Genomics* 88: 779-790.
- Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610823. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Robinson, A.J., Overy, C. and Kunji, E.R. 2008. The mechanism of transport by mitochondrial carriers based on analysis of symmetry. *Proc. Natl. Acad. Sci. USA* 105: 17766-17771.
- Fiermonte, G., Paradies, E., Todisco, S., Marobbio, C.M. and Palmieri, F. 2009. A novel member of solute carrier family 25 (SLC25A42) is a transporter of coenzyme A and adenosine 3',5'-diphosphate in human mitochondria. *J. Biol. Chem.* 284: 18152-18159.
- Iacopetta, D., Carrisi, C., De Filippis, G., Calcagnile, V.M., Cappello, A.R., Chimento, A., Curcio, R., Santoro, A., Voza, A., Dolce, V., Palmieri, F. and Capobianco, L. 2010. The biochemical properties of the mitochondrial thiamine pyrophosphate carrier from *Drosophila melanogaster*. *FEBS J.* 277: 1172-1181.

CHROMOSOMAL LOCATION

Genetic locus: SLC25A42 (human) mapping to 19p13.11; Slc25a42 (mouse) mapping to 8 B3.3.

SOURCE

SLC25A42 (T-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of SLC25A42 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244163 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SLC25A42 (T-16) is recommended for detection of SLC25A42 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SLC25A42 (T-16) is also recommended for detection of SLC25A42 in additional species, including canine.

Suitable for use as control antibody for SLC25A42 siRNA (h): sc-97761, SLC25A42 siRNA (m): sc-153518, SLC25A42 shRNA Plasmid (h): sc-97761-SH, SLC25A42 shRNA Plasmid (m): sc-153518-SH, SLC25A42 shRNA (h) Lentiviral Particles: sc-97761-V and SLC25A42 shRNA (m) Lentiviral Particles: sc-153518-V.

Molecular Weight of SLC25A42: 35 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.