

SPACA5 (D-15): sc-244196

BACKGROUND

SPACA5 (sperm acrosome-associated protein 5), also known as lysozyme-like protein 5 or sperm-specific lysozyme-like protein X (SLLP-X), is a 150 amino acid secreted protein belonging to the glycosyl hydrolase 22 family. Acting as a component of the perinuclear theca and the acrosome, two structures present on the sperm head, SPACA5 may be a factor in sperm head defects. The gene encoding SPACA5 maps to human chromosome Xp11.23. Chromosome X consists of about 153 million base pairs and nearly 1,000 genes. Color blindness, hemophilia, and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently than females as males carry a single X chromosome.

REFERENCES

1. Cramer, E.M. and Breton-Gorius, J. 1987. Ultrastructural localization of lysozyme in human neutrophils by immunogold. *J. Leukoc. Biol.* 41: 242-247.
2. Zhang, K., Gao, R., Zhang, H., Cai, X., Shen, C., Wu, C., Zhao, S. and Yu, L. 2005. Molecular cloning and characterization of three novel lysozyme-like genes, predominantly expressed in the male reproductive system of humans, belonging to the c-type lysozyme/ α -lactalbumin family. *Biol. Reprod.* 73: 1064-1071.
3. Augui, S., Filion, G.J., Huart, S., Nora, E., Guggiari, M., Maresca, M., Stewart, A.F. and Heard, E. 2007. Sensing X chromosome pairs before X inactivation via a novel X-pairing region of the Xic. *Science* 318: 1632-1636.
4. Shustrova, E.N. and Artamonova, I.I. 2009. Evolutionary history of the SSX family of human C/T-antigens. *Mol. Biol.* 43: 1025-1031.
5. Cocquet, J., Ellis, P.J., Yamauchi, Y., Mahadevaiah, S.K., Affara, N.A., Ward, M.A. and Burgoyne, P.S. 2009. The multicopy gene Sly represses the sex chromosomes in the male mouse germline after meiosis. *PLoS Biol.* 7: e1000244.
6. Leahy, T., Marti, J.I., Crossett, B., Evan, G. and Maxwell, W.M. 2011. Two-dimensional polyacrylamide gel electrophoresis of membrane proteins from flow cytometrically sorted ram sperm. *Theriogenology* 75: 962-971.

CHROMOSOMAL LOCATION

Genetic locus: SPACA5 (human) mapping to Xp11.23; Spaca5 (mouse) mapping to X A1.3.

SOURCE

SPACA5 (D-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of SPACA5 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244196 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SPACA5 (D-15) is recommended for detection of SPACA5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with SPACA1, SPACA3 or SPACA4.

SPACA5 (D-15) is also recommended for detection of SPACA5 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for SPACA5 siRNA (h): sc-91003, SPACA5 siRNA (m): sc-153698, SPACA5 shRNA Plasmid (h): sc-91003-SH, SPACA5 shRNA Plasmid (m): sc-153698-SH, SPACA5 shRNA (h) Lentiviral Particles: sc-91003-V and SPACA5 shRNA (m) Lentiviral Particles: sc-153698-V.

Molecular Weight of SPACA5: 18 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.