

SPINK9 (K-16): sc-244215

BACKGROUND

SPINK9 (serine peptidase inhibitor, Kazal type 9), also known as LEKTI2, is an 86 amino acid secreted protein that contains one kazal-like domain and is thought to function as a serine protease inhibitor, possibly playing a role in proteolytic cascades. The gene encoding SPINK9 maps to human chromosome 5, which contains 181 million base pairs and comprises nearly 6% of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

- Dixon, M.J., Read, A.P., Donnai, D., Colley, A., Dixon, J. and Williamson, R. 1991. The gene for Treacher Collins syndrome maps to the long arm of chromosome 5. *Am. J. Hum. Genet.* 49: 17-22.
- Joslyn, G., Carlson, M., Thliveris, A., Albertsen, H., Gelbert, L., Samowitz, W., Groden, J., Stevens, J., Spirio, L. and Robertson, M. 1991. Identification of deletion mutations and three new genes at the familial polyposis locus. *Cell* 66: 601-613.
- Kinzler, K.W., Nilbert, M.C., Su, L.K., Vogelstein, B., Bryan, T.M., Levy, D.B., Smith, K.J., Preisinger, A.C., Hedge, P. and McKechnie, D. 1991. Identification of FAP locus genes from chromosome 5q21. *Science* 253: 661-665.
- Nishisho, I., Nakamura, Y., Miyoshi, Y., Miki, Y., Ando, H., Horii, A., Koyama, K., Utsunomiya, J., Baba, S. and Hedge, P. 1991. Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science* 253: 665-669.
- Prieschl, E.E., Pendl, G.G., Harrer, N.E. and Baumruker, T. 1996. The murine homolog of TB2/DP1, a gene of the familial adenomatous polyposis (FAP) locus. *Gene* 169: 215-218.
- Shin, S.M., Chung, Y.J., Oh, S.T., Jeon, H.M., Hwang, L.J., Namkoong, H., Kim, H.K., Cho, G.W., Hur, S.Y., Kim, T.E., Lee, Y.S., Park, Y.G., Ko, J. and Kim, J.W. 2006. HCCR-1-interacting molecule "deleted in polyposis 1" plays a tumor-suppressor role in colon carcinogenesis. *Gastroenterology* 130: 2074-2086.
- Brattsand, M., Stefansson, K., Hubiche, T., Nilsson, S.K. and Egelrud, T. 2009. SPINK9: SPINK9: a selective, skin-specific Kazal-type serine protease inhibitor. *J. Invest. Dermatol.* 129: 1656-1665.
- Meyer-Hoffert, U., Wu, Z. and Schröder, J.M. 2009. Identification of lympho-epithelial Kazal-type inhibitor 2 in human skin as a kallikrein-related peptidase 5-specific protease inhibitor. *PLoS ONE* 4: e4372.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: SPINK9 (human) mapping to 5q32.

SOURCE

SPINK9 (K-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of SPINK9 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244215 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SPINK9 (K-16) is recommended for detection of SPINK9 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other SPINK family members.

Suitable for use as control antibody for SPINK9 siRNA (h): sc-91686, SPINK9 shRNA Plasmid (h): sc-91686-SH and SPINK9 shRNA (h) Lentiviral Particles: sc-91686-V.

Molecular Weight of SPINK9: 10 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.