

Tom6 (C-14): sc-244477

BACKGROUND

The mitochondrial preprotein translocases of the outer membrane (Tom) form a multisubunit complex that facilitates the import of nuclear-encoded precursor proteins across the mitochondrial outer membrane. The Tom machinery consists of import receptors for the initial binding of cytosolically synthesized preproteins, and a general import pore (GIP) for the membrane translocation of various preproteins into the mitochondrion. Tom6 (translocase of outer mitochondrial membrane 6 homolog (yeast)), also known as OBTP, is a 74 amino acid protein that belongs to the Tom6 family. Localizing to the mitochondrial outer membrane, Tom6 forms part of the preprotein translocase complex of the outer mitochondrial membrane (TOM complex). Tom6 is encoded by a gene located on human chromosome 6p25.3 and mouse chromosome 17 C.

REFERENCES

- Rapaport, D., Neupert, W. and Lill, R. 1997. Mitochondrial protein import. Tom40 plays a major role in targeting and translocation of preproteins by forming a specific binding site for the presequence. *J. Biol. Chem.* 272: 18725-18731.
- Dekker, P.J., Ryan, M.T., Brix, J., Muller, H., Honlinger, A. and Pfanner, N. 1998. Preprotein translocase of the outer mitochondrial membrane: molecular dissection and assembly of the general import pore complex. *Mol. Cell. Biol.* 18: 6515-6524.
- Yano, M., Kanazawa, M., Terada, K., Takeya, M., Hoogenraad, N. and Mori, M. 1998. Functional analysis of human mitochondrial receptor Tom20 for protein import into mitochondria. *J. Biol. Chem.* 273: 26844-26851.
- Rapaport, D. and Neupert, W. 1999. Biogenesis of Tom40, core component of the TOM complex of mitochondria. *J. Cell Biol.* 146: 321-331.
- Ahting, U., Thun, C., Hegerl, R., Typke, D., Nargang, F.E., Neupert, W. and Nussberger, S. 1999. The TOM core complex: the general protein import pore of the outer membrane of mitochondria. *J. Cell Biol.* 147: 959-968.
- Ryan, M.T., Wagner, R. and Pfanner, N. 2000. The transport machinery for the import of preproteins across the outer mitochondrial membrane. *Int. J. Biochem. Cell Biol.* 32: 13-21.
- Kato, H. and Mihara, K. 2008. Identification of Tom5 and Tom6 in the preprotein translocase complex of human mitochondrial outer membrane. *Biochem. Biophys. Res. Commun.* 369: 958-963.
- Dukanovic, J., Dimmer, K.S., Bonnefoy, N., Krumpke, K. and Rapaport, D. 2009. Genetic and functional interactions between the mitochondrial outer membrane proteins Tom6 and Sam37. *Mol. Cell. Biol.* 29: 5975-5988.

CHROMOSOMAL LOCATION

Genetic locus: TOMM6 (human) mapping to 6p25.3; Tomm6 (mouse) mapping to 17 C.

SOURCE

Tom6 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of Tom6 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244477 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Tom6 (C-14) is recommended for detection of Tom6 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Tom family members.

Tom6 (C-14) is also recommended for detection of Tom6 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for Tom6 siRNA (m): sc-152464, Tom6 shRNA Plasmid (m): sc-152464-SH and Tom6 shRNA (m) Lentiviral Particles: sc-152464-V.

Molecular Weight of Tom6: 8 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.