

# ZCCHC4 (C-12): sc-244710

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZCCHC4 (zinc finger, CCHC domain containing 4) is a 513 amino acid protein that contains one CCHC-type zinc finger and one DHHC-type zinc finger and is thought to function as a methyltransferase. The gene encoding ZCCHC4 maps to human chromosome 4, which encodes nearly 6% of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes. Defects in some of the genes located on chromosome 4 are associated with Huntington's disease, Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

## REFERENCES

- Albrecht, M. and Lengauer, T. 2004. Novel Sm-like proteins with long C-terminal tails and associated methyltransferases. *FEBS Lett.* 569: 18-26.
- Cowan, C.M. and Raymond, L.A. 2006. Selective neuronal degeneration in Huntington's disease. *Curr. Top. Dev. Biol.* 75:25-71.
- Cunningham, M.L., Seto, M.L., Ratisoontorn, C., Heike, C.L. and Hing, A.V. 2007. Syndromic craniosynostosis: from history to hydrogen bonds. *Orthod. Craniofac. Res.* 10: 67-81.
- Doherty, E.S., Lacbawan, F., Hadley, D.W., Brewer, C., Zalewski, C., Kim, H.J., Solomon, B., Rosenbaum, K., Domingo, D.L., Hart, T.C., Brooks, B.P., Immken, L., Lowry, R.B., Kimonis, V., Shanske, A.L., Jehee, F.S., Bueno, M.R., Knightly, C., McDonald-McGinn, D., Zackai, E.H. and Muenke, M. 2007. Muenke syndrome (FGFR3-related craniosynostosis): Expansion of the phenotype and review of the literature. *Am. J. Med. Genet. A* 143: 3204-3215.
- Ruiz-Perez, V.L., Blair, H.J., Rodriguez-Andres, M.E., Blanco, M.J., Wilson, A., Liu, Y.N., Miles, C., Peters, H. and Goodship, J.A. 2007. Evc is a positive mediator of Ihh-regulated bone growth that localises at the base of chondrocyte cilia. *Development* 134: 2903-2912.
- Stack, E.C., Dedeoglu, A., Smith, K.M., Cormier, K., Kubilus, J.K., Bogdanov, M., Matson, W.R., Yang, L., Jenkins, B.G., Luthi-Carter, R., Kowall, N.W., Hersch, S.M., Beal, M.F. and Ferrante, R.J. 2007. Neuroprotective effects of synaptic modulation in Huntington's disease R6/2 mice. *J. Neurosci.* 27: 12908-12915.
- Versteegh, F.G., Buma, S.A., Costin, G., de Jong, W.C. and Hennekam, R.C. 2007. Evc Working Party. Growth hormone analysis and treatment in Ellis-van Creveld syndrome. *Am. J. Med. Genet. A* 143: 2113-2121.
- Kang, B., Hao, C., Wang, H., Zhang, J., Xing, R., Shao, J., Li, W., Xu, N., Lu, Y. and Liu, S. 2008. Evaluation of hepatic-metastasis risk of colorectal cancer upon the protein signature of PI3K/AKT pathway. *J. Proteome Res.* 7: 3507-3515.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## CHROMOSOMAL LOCATION

Genetic locus: ZCCHC4 (human) mapping to 4p15.2.

## SOURCE

ZCCHC4 (C-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of ZCCHC4 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244710 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

ZCCHC4 (C-12) is recommended for detection of ZCCHC4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of ZCCHC4: 59 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.