

ZNF618 (D-14): sc-244814

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF618 (zinc finger protein 618), also known as NEDD10, is a 954 amino acid nuclear protein that belongs to the Krüppel C₂H₂-type zinc-finger protein family and may play a role in transcriptional regulation. Containing four C₂H₂-type zinc fingers, ZNF618 exists as three alternatively spliced isoforms that are encoded by a gene located on human chromosome 9q32.

REFERENCES

1. Witzgall, R., O'Leary, E., Leaf, A., Onaldi, D. and Bonventre, J.V. 1994. The Krüppel-associated box-A (KRAB-A) domain of zinc finger proteins mediates transcriptional repression. *Proc. Natl. Acad. Sci. USA* 91: 4514-4518.
2. Mark, C., Abrink, M. and Hellman, L. 1999. Comparative analysis of KRAB zinc finger proteins in rodents and man: evidence for several evolutionarily distinct subfamilies of KRAB zinc finger genes. *DNA Cell Biol.* 18: 381-396.
3. Williams, A.J., Blacklow, S.C. and Collins, T. 1999. The zinc finger-associated SCAN box is a conserved oligomerization domain. *Mol. Cell. Biol.* 19: 8526-8535.
4. Peng, H., Begg, G.E., Harper, S.L., Friedman, J.R., Speicher, D.W. and Rauscher, F.J. 2000. Biochemical analysis of the Krüppel-associated box (KRAB) transcriptional repression domain. *J. Biol. Chem.* 275: 18000-18010.
5. Urrutia, R. 2003. KRAB-containing zinc-finger repressor proteins. *Genome Biol.* 4: 231.
6. Huntley, S., Baggott, D.M., Hamilton, A.T., Tran-Gyamfi, M., Yang, S., Kim, J., Gordon, L., Branscomb, E. and Stubbs, L. 2006. A comprehensive catalog of human KRAB-associated zinc finger genes: insights into the evolutionary history of a large family of transcriptional repressors. *Genome Res.* 16: 669-677.
7. Tian, C.Y., Zhang, L.Q. and He, F.C. 2006. Progress in the study of KRAB zinc finger protein. *Yi Chuan* 28: 1451-1456.

CHROMOSOMAL LOCATION

Genetic locus: ZNF618 (human) mapping to 9q32; Zfp618 (mouse) mapping to 4 B3.

SOURCE

ZNF618 (D-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ZNF618 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-244814 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ZNF618 (D-14) is recommended for detection of ZNF618 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZNF family members.

ZNF618 (D-14) is also recommended for detection of ZNF618 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for ZNF618 siRNA (h): sc-92635, ZNF618 siRNA (m): sc-155764, ZNF618 shRNA Plasmid (h): sc-92635-SH, ZNF618 shRNA Plasmid (m): sc-155764-SH, ZNF618 shRNA (h) Lentiviral Particles: sc-92635-V and ZNF618 shRNA (m) Lentiviral Particles: sc-155764-V.

Molecular Weight of ZNF618 isoforms: 105/96/31 kDa.

Positive Controls: NTERA-2 cl.D1 whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotting A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.