

BTNL8 (D-17): sc-245053

BACKGROUND

BTNL8 (butyrophilin-like 8) is a 500 amino acid single-pass type I membrane protein that belongs to the immunoglobulin superfamily and contains one B30.2/SPRY domain and one Ig-like V-type (immunoglobulin-like) domain. Expressed as multiple alternatively spliced isoforms, BTNL8 is encoded by a gene which maps to human chromosome 5. With 181 million base pairs encoding around 1,000 genes, chromosome 5 comprises about 6% of human genomic DNA. Deletion of the p arm of chromosome 5 leads to cri du chat syndrome, while deletion of the q arm on chromosome 5 is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

- Dixon, M.J., et al. 1991. The gene for Treacher Collins syndrome maps to the long arm of chromosome 5. *Am. J. Hum. Genet.* 49: 17-22.
- Joslyn, G., et al. 1991. Identification of deletion mutations and three new genes at the familial polyposis locus. *Cell* 66: 601-613.
- Kinzler, K.W., et al. 1991. Identification of FAP locus genes from chromosome 5q21. *Science* 253: 661-665.
- Nishisho, I., et al. 1991. Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science* 253: 665-669.
- Prieschl, E.E., et al. 1996. The murine homolog of TB2/DP1, a gene of the familial adenomatous polyposis (FAP) locus. *Gene* 169: 215-218.
- Puente, X.S., et al. 2004. A genomic analysis of rat proteases and protease inhibitors. *Genome Res.* 14: 609-622.
- Shin, S.M., et al. 2006. HCCR-1-interacting molecule "deleted in polyposis 1" plays a tumor-suppressor role in colon carcinogenesis. *Gastroenterology* 130: 2074-2086.

CHROMOSOMAL LOCATION

Genetic locus: BTNL8 (human) mapping to 5q35.3.

SOURCE

BTNL8 (D-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of BTNL8 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-245053 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

BTNL8 (D-17) is recommended for detection of BTNL8 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with BTNL2, BTNL3, or 9.


Suitable for use as control antibody for BTNL8 siRNA (h): sc-91741, BTNL8 shRNA Plasmid (h): sc-91741-SH and BTNL8 shRNA (h) Lentiviral Particles: sc-91741-V.

Molecular Weight of BTNL8: 57 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



BTNL8 (D-17): sc-245053. Immunofluorescence staining of formalin-fixed HepG2 cells showing membrane localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.