C4orf19 (S-20): sc-245748



The Power to Question

BACKGROUND

C4orf19 (chromosome 4 open reading frame 12) is a 314 amino acid protein encoded by a gene that maps to human chromosome 4. The mouse homolog, 0610040J01Rik, is a 313 amino acid protein encoded by a gene mapping to mouse chromosome 5. Human chromosome 4 represents approximately 6% of the human genome and contains nearly 900 genes. Notably, the Huntingtin gene, which is found to encode an expanded glutamine tract in cases of Huntington's disease, is on chromosome 4. FGFR-3 is also encoded by a gene that is located on chromosome 4, and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also tied to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

REFERENCES

- 1. Hillier, L.W., et al. 2005. Generation and annotation of the DNA sequences of human chromosomes 2 and 4. Nature 434: 724-731.
- 2. Cowan, C.M. and Raymond, L.A. 2006. Selective neuronal degeneration in Huntington's disease. Curr. Top. Dev. Biol. 75:25-71.
- 3. Chandler, R.J., et al. 2007. Metabolic phenotype of methylmalonic acidemia in mice and humans: the role of skeletal muscle. BMC Med. Genet. 8: 64.
- 4. Cunningham, M.L., et al. 2007. Syndromic craniosynostosis: from history to hydrogen bonds. Orthod. Craniofac. Res. 10: 67-81.
- de Frutos, C.A., et al. 2007. Snail1 is a transcriptional effector of FGFR3 signaling during chondrogenesis and achondroplasias. Dev. Cell 13: 872-883.
- 6. Versteegh, F.G., et al. 2007. Growth hormone analysis and treatment in Ellis-van Creveld syndrome. Am. J. Med. Genet. A 143A: 2113-2121.
- Doherty, E.S., et al. 2007. Muenke syndrome (FGFR3-related craniosynostosis): Expansion of the phenotype and review of the literature. Am. J. Med. Genet. A 143A: 3204-3215.
- 8. Ruiz-Perez, V.L., et al. 2007. Evc is a positive mediator of lhh-regulated bone growth that localises at the base of chondrocyte cilia. Development 134: 2903-2912.
- 9. Stack, E.C., et al. 2007. Neuroprotective effects of synaptic modulation in Huntington's disease R6/2 mice. J. Neurosci. 27: 12908-12915.

CHROMOSOMAL LOCATION

Genetic locus: C4orf19 (human) mapping to 4p14.

SOURCE

C4orf19 (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of C4orf19 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-245748 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

C4orf19 (S-20) is recommended for detection of C4orf19 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for C4orf19 siRNA (h): sc-89176, C4orf19 shRNA Plasmid (h): sc-89176-SH and C4orf19 shRNA (h) Lentiviral Particles: sc-89176-V.

Molecular Weight of C4orf19: 34 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**