

# CCNI2 (K-16): sc-246217

## BACKGROUND

CCNI2 (cyclin-I2) is a 369 amino acid protein that belongs to the cyclin family. The gene encoding CCNI2 consists of approximately 6,720 bases and maps to human chromosome 5q31.1. Containing 181 million base pairs, chromosome 5 comprises nearly 6% of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

## REFERENCES

1. Edwards, S.J., Gladwin, A.J. and Dixon, M.J. 1997. The mutational spectrum in Treacher Collins syndrome reveals a predominance of mutations that create a premature-termination codon. *Am. J. Hum. Genet.* 60: 515-524.
2. McDaniel, L.D., Legerski, R., Lehmann, A.R., Friedberg, E.C. and Schultz, R.A. 1997. Confirmation of homozygosity for a single nucleotide substitution mutation in a Cockayne syndrome patient using monoallelic mutation analysis in somatic cell hybrids. *Hum. Mutat.* 10: 317-321.
3. Crawford, M.J., Lanctôt, C., Tremblay, J.J., Jenkins, N., Gilbert, D., Copeland, N., Beatty, B. and Drouin, J. 1997. Human and murine PTX1/Ptx1 gene maps to the region for Treacher Collins syndrome. *Mamm. Genome* 8: 841-845.
4. Finch, R., Moore, H.G., Lindor, N., Jalal, S.M., Markowitz, A., Suresh, J., Offit, K. and Guillem, J.G. 2005. Familial adenomatous polyposis and mental retardation caused by a *de novo* chromosomal deletion at 5q15-q22: report of a case. *Dis. Colon Rectum* 48: 2148-2152.
5. Anindya, R., Aygün, O. and Svejstrup, J.Q. 2007. Damage-induced ubiquitylation of human RNA polymerase II by the ubiquitin ligase Nedd4, but not Cockayne syndrome proteins or BRCA1. *Mol. Cell* 28: 386-397.
6. Vera-Carbonell, A., Bafalliu, J.A., Guillen-Navarro, E., Escalona, A., Ballesta-Martínez, M.J., Fuster, C., Fernández, A. and López-Expósito, I. 2009. Characterization of a *de novo* complex chromosomal rearrangement in a patient with cri-du-chat and trisomy 5p syndromes. *Am. J. Med. Genet. A* 149A: 2513-2521.
7. Ravandi, F., Issa, J.P., Garcia-Manero, G., O'Brien, S., Pierce, S., Shan, J., Borthakur, G., Verstovsek, S., Faderl, S., Cortes, J. and Kantarjian, H. 2009. Superior outcome with hypomethylating therapy in patients with acute myeloid leukemia and high-risk myelodysplastic syndrome and chromosome 5 and 7 abnormalities. *Cancer* 115: 5746-5751.
8. Sazawal, S., Kumar, B., Hasan, S.K., Dutta, P., Kumar, R., Chaubey, R., Mir, R. and Saxena, R. 2009. Haematological & molecular profile of acute myelogenous leukaemia in India. *Indian J. Med. Res.* 129: 256-261.

## CHROMOSOMAL LOCATION

Genetic locus: CCNI2 (human) mapping to 5q31.1.

## SOURCE

CCNI2 (K-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CCNI2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-246217 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

CCNI2 (K-16) is recommended for detection of CCNI2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CCNI2 (K-16) is also recommended for detection of CCNI2 in additional species, including bovine.

Suitable for use as control antibody for CCNI2 siRNA (h): sc-91752, CCNI2 shRNA Plasmid (h): sc-91752-SH and CCNI2 shRNA (h) Lentiviral Particles: sc-91752-V.

Molecular Weight of CCNI2: 41 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.