# SANTA CRUZ BIOTECHNOLOGY, INC.

# DENND2A (P-19): sc-246427



## BACKGROUND

DENND2A (DENN/MADD domain containing 2A), also known as KIAA1277, is a 1,009 amino acid protein that contains a dDENN domain, DENN domain, and uDENN domain and exists as 2 isoforms as a result of alternative splicing. The DENND2A protein is thought to target to actin filaments and control Rab9-dependent trafficking of mannose-6-phosphate receptor to lysosomes. The gene encoding DENND2A maps to human chromosome 7, which encodes over 1,000 genes and makes up about 5% of the human genome. Chromosome 7 has been linked to osteogenesis imperfecta, Pendred syndrome, lissencephaly, citrullinemia and Shwachman-Diamond syndrome. The deletion of a portion of the q arm of chromosome 7 is associated with Williams-Beuren syndrome, a condition characterized by mild mental retardation, an unusual comfort and friendliness with strangers and an elfin appearance. Deletions of portions of the q arm of chromosome 7 are also seen in a number of myeloid disorders including cases of acute myelogenous leukemia and myelodysplasia.

## REFERENCES

- 1. Tsipouras, P., et al. 1983. Restriction fragment length polymorphism associated with the pro  $\alpha$  2(I) gene of human type I procollagen. Application to a family with an autosomal dominant form of osteogenesis imperfecta. J. Clin. Invest. 72: 1262-1267.
- Liang, H., et al. 1998. Molecular anatomy of chromosome 7q deletions in myeloid neoplasms: evidence for multiple critical loci. Proc. Natl. Acad. Sci. USA 95: 3781-3785.
- 3. Hillier, L.W., et al. 2003. The DNA sequence of human chromosome 7. Nature 424: 157-164.
- Eckert, M.A., et al. 2006. The neurobiology of Williams syndrome: cascading influences of visual system impairment? Cell. Mol. Life Sci. 63: 1867-1875.
- Osborne, L.R., et al. 2006. Williams-Beuren syndrome diagnosis using fluorescence *in situ* hybridization. Methods Mol. Med. 126: 113-128.
- Reiner, O., et al. 2006. Lissencephaly 1 linking to multiple diseases: mental retardation, neurodegeneration, schizophrenia, male sterility, and more. Neuromolecular Med. 8: 547-565.
- Shimamura, A. 2006. Shwachman-Diamond syndrome. Semin. Hematol. 43: 178-188.

## CHROMOSOMAL LOCATION

Genetic locus: Dennd2a (mouse) mapping to 6 B1.

#### SOURCE

DENND2A (P-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of DENND2A of mouse origin.

## PRODUCT

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-246427 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

DENND2A (P-19) is recommended for detection of DENND2A of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with DENND2C or DENND2D.

Suitable for use as control antibody for DENND2A siRNA (m): sc-142998, DENND2A shRNA Plasmid (m): sc-142998-SH and DENND2A shRNA (m) Lentiviral Particles: sc-142998-V.


Molecular Weight of DENND2A: 114/90 kDa.

Positive Controls: mouse brain extract: sc-2253.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### DATA



DENND2A (P-19): sc-246427. Western blot analysis of DENND2A expression in mouse brain tissue extract.

#### **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.