# SANTA CRUZ BIOTECHNOLOGY, INC.

# FAM108C1 (P-13): sc-246545



BACKGROUND

FAM108C1 is a 329 amino acid protein that exists as 2 alternatively spliced isoforms. The gene encoding FAM108C1 maps to human chromosome 15, which encodes more than 700 genes and is made up of approximately 106 million base pairs. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. In the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13 encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

## REFERENCES

- Cachón-González, M.B., et al. 2006. Effective gene therapy in an authentic model of Tay-Sachs-related diseases. Proc. Natl. Acad. Sci. USA 103: 10373-10378.
- 2. Zody, M.C., et al. 2006. Analysis of the DNA sequence and duplication history of human chromosome 15. Nature 440: 671-675.
- 3. Diene, G., et al. 2007. The Prader-Willi syndrome. Ann. Endocrinol. 68: 129-137.
- 4. Lalande, M., et al. 2007. Molecular epigenetics of Angelman syndrome. Cell. Mol. Life Sci. 64: 947-960.
- Maegawa, G.H., et al. 2007. Pyrimethamine as a potential pharmacological chaperone for late-onset forms of GM2 gangliosidosis. J. Biol. Chem. 282: 9150-9161.
- Makoff, A.J., et al. 2007. Detailed analysis of 15q11-q14 sequence corrects errors and gaps in the public access sequence to fully reveal large segmental duplications at breakpoints for Prader-Willi, Angelman, and inv dup(15) syndromes. Genome Biol. 8: R114.
- 7. Ramirez, F., et al. 2007. Fibrillin-rich microfibrils: Structural determinants of morphogenetic and homeostatic events. J. Cell. Physiol. 213: 326-330.

# CHROMOSOMAL LOCATION

Genetic locus: FAM108C1 (human) mapping to 15q25.1; Fam108c (mouse) mapping to 7 D3.

#### SOURCE

FAM108C1 (P-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FAM108C1 of human origin.

# PRODUCT

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-246545 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### APPLICATIONS

FAM108C1 (P-13) is recommended for detection of FAM108C1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other FAM108 family members.

FAM108C1 (P-13) is also recommended for detection of FAM108C1 in additional species, including bovine and porcine.

Suitable for use as control antibody for FAM108C1 siRNA (h): sc-90123, FAM108C1 siRNA (m): sc-108631, FAM108C1 shRNA Plasmid (h): sc-90123-SH, FAM108C1 shRNA Plasmid (m): sc-108631-SH, FAM108C1 shRNA (h) Lentiviral Particles: sc-90123-V and FAM108C1 shRNA (m) Lentiviral Particles: sc-108631-V.

Molecular Weight of FAM108C1: 36 kDa.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.