

GDPD3 (V-17): sc-246991

BACKGROUND

Phosphodiesterases (PDEs) are important for the downregulation of the intracellular level of the second messenger cyclic adenosine monophosphate (cAMP) by hydrolyzing cAMP to 5'AMP. The PDE family contains proteins that serve tissue-specific roles in the regulation of lipolysis, glycogenolysis, myocardial contractility and smooth muscle relaxation. GDPD3 (glycerophosphodiester phosphodiesterase domain-containing protein 3) is a 318 amino acid multi-pass membrane protein that belongs to the glycerophosphoryl diester phosphodiesterase family and contains one GDPD domain. Existing as two alternatively spliced isoforms, GDPD3 is encoded by a gene that maps to human chromosome 16p11.2. Chromosome 16 encodes over 900 genes and comprises nearly 3% of the human genome. Giant axonal neuropathy, Rubinstein-Taybi syndrome and Crohn's disease are associated with defects in chromosome 16.

REFERENCES

1. Baraitser, M., et al. 1983. The Rubinstein-Taybi syndrome: occurrence in two sets of identical twins. *Clin. Genet.* 23: 318-320.
2. Breuning, M.H., et al. 1993. Rubinstein-Taybi syndrome caused by submicroscopic deletions within 16p13.3. *Am. J. Hum. Genet.* 52: 249-254.
3. Bomont, P., et al. 2000. The gene encoding gigaxonin, a new member of the cytoskeletal BTB/kelch repeat family, is mutated in giant axonal neuropathy. *Nat. Genet.* 26: 370-374.
4. Kuhlensäumer, G., et al. 2002. Giant axonal neuropathy (GAN): case report and two novel mutations in the gigaxonin gene. *Neurology* 58: 1273-1276.
5. Cho, J.H. 2004. Advances in the genetics of inflammatory bowel disease. *Curr. Gastroenterol. Rep.* 6: 467-473.
6. Mathew, C.G., et al. 2004. Genetics of inflammatory bowel disease: progress and prospects. *Hum. Mol. Genet.* 13: R161-R168.
7. Martin, J., et al. 2004. The sequence and analysis of duplication-rich human chromosome 16. *Nature* 432: 988-994.

CHROMOSOMAL LOCATION

Genetic locus: GDPD3 (human) mapping to 16p11.2; Gdpd3 (mouse) mapping to 7 F3.

SOURCE

GDPD3 (V-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of GDPD3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-246991 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

GDPD3 (V-17) is recommended for detection of GDPD3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GDPD3 (V-17) is also recommended for detection of GDPD3 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for GDPD3 siRNA (h): sc-93057, GDPD3 siRNA (m): sc-145377, GDPD3 shRNA Plasmid (h): sc-93057-SH, GDPD3 shRNA Plasmid (m): sc-145377-SH, GDPD3 shRNA (h) Lentiviral Particles: sc-93057-V and GDPD3 shRNA (m) Lentiviral Particles: sc-145377-V.


Molecular Weight of GDPD3 isoforms: 37/30 kDa.

Positive Controls: mouse liver extract: sc-2256.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GDPD3 (V-17): sc-246991. Western blot analysis of GDPD3 expression in mouse liver tissue extract.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.