KIAA1731 (S-16): sc-247334



The Power to Question

BACKGROUND

KIAA1731 is a 2,601 amino acid protein that is implicated in centriole formation or stability. Existing as four alternatively spliced isoforms, KIAA1731 is encoded by a gene that maps to human chromosome 11q21. Chromosome 11 comprises approximately 4% of human genomic DNA and is considered a gene and disease association dense chromosome. The chromosome 11 encoded Atm gene is important for regulation of cell cycle arrest and apoptosis following double strand DNA breaks. Atm mutation leads to the disorder known as ataxia-telangiectasia. The blood disorders Sickle cell anemia and thalassemia are caused by HBB gene mutations, while Wilms' tumors, WAGR syndrome and Denys-Drash syndrome are associated with mutations of the WT1 gene. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are also associated with defects in chromosome 11-encoded genes.

REFERENCES

- 1. Fabiani, J.E., et al. 2000. Hereditary angioedema. Long-term follow-up of 88 patients. Experience of the Argentine allergy and immunology institute. Allergol. Immunopathol. 28: 267-271.
- Jira, P.E., et al. 2003. Smith-lemli-opitz syndrome and the DHCR7 gene. Ann. Hum. Genet. 67: 269-280.
- Schuchman, E.H. 2007. The pathogenesis and treatment of acid sphingomyelinase-deficient Niemann-pick disease. J. Inherit. Metab. Dis. 30: 654-663
- Siem, G., et al. 2008. Jervell and Lange-nielsen syndrome in Norwegian children: aspects around cochlear implantation, hearing, and balance. Ear Hear. 29: 261-269.
- Bhuiyan, Z.A., et al. 2008. An intronic mutation leading to incomplete skipping of exon-2 in KCNQ1 rescues hearing in Jervell and Lange-nielsen syndrome. Prog. Biophys. Mol. Biol. 98: 319-327.
- Coldren, C.D., et al. 2009. Chromosomal microarray mapping suggests a role for BSX and Neurogranin in neurocognitive and behavioral defects in the 11q terminal deletion disorder (Jacobsen syndrome). Neurogenetics 10: 89-95.

CHROMOSOMAL LOCATION

Genetic locus: KIAA1731 (human) mapping to 11q21; 5830418K08Rik (mouse) mapping to 9 A2.

SOURCE

KIAA1731 (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of KIAA1731 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-247334 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

KIAA1731 (S-16) is recommended for detection of KIAA1731 of human origin, 5830418K08Rik of mouse origin and the corresponding rat homolog by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 5830418K08Rik siRNA (m): sc-140406, 5830418K08Rik shRNA Plasmid (m): sc-140406-SH and 5830418K08Rik shRNA (m) Lentiviral Particles: sc-140406-V.

Molecular Weight of KIAA1731 isoforms: 295/88/295/69 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**