

LANCL3 (D-17): sc-247406

BACKGROUND

LANCL3 (LanC lantibiotic synthetase component C-like 3) is a 420 amino acid protein that belongs to the LanC-like protein family and exists as 2 alternatively spliced isoforms. The gene encoding LANCL3 maps to human chromosome Xp21.1 and mouse chromosome X A1.1. The X and Y chromosomes are the human sex chromosomes. Chromosome X consists of about 153 million base pairs and nearly 1,000 genes. The combination of an X and Y chromosome leads to normal male development while two copies of X leads to normal female development. There are a number of conditions related to an unusual number and combination of sex chromosomes being inherited, including Turner's syndrome, Klinefelter's syndrome and Triple X syndrome. Color blindness, hemophilia, and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently as males carry a single X chromosome.

REFERENCES

- Gubin, A.N., et al. 1999. Gene expression in proliferating human erythroid cells. *Genomics* 59: 168-177.
- Gianfrancesco, F., et al. 2001. Differential divergence of three human pseudoautosomal genes and their mouse homologs: implications for sex chromosome evolution. *Genome Res.* 11: 2095-2100.
- Bernardino-Sgherri, J., et al. 2002. Overall DNA methylation and chromatin structure of normal and abnormal X chromosomes. *Cytogenet. Genome Res.* 99: 85-91.
- Deeb, S.S. 2005. The molecular basis of variation in human color vision. *Clin. Genet.* 67: 369-377.
- Bojesen, A., et al. 2006. The metabolic syndrome is frequent in Klinefelter's syndrome and is associated with abdominal obesity and hypogonadism. *Diabetes Care* 29: 1591-1598.
- Maggio, M.C., et al. 2007. Polycystic ovary and gonadoblastoma in Turner's syndrome. *Minerva Pediatr.* 59: 397-401.
- Helderman-van den Enden, A.T., et al. 2009. Recurrence risk due to germ line mosaicism: duchenne and becker muscular dystrophy. *Clin. Genet.* 75: 465-472.

CHROMOSOMAL LOCATION

Genetic locus: LANCL3 (human) mapping to Xp21.1; Lancl3 (mouse) mapping to X A1.1.

SOURCE

LANCL3 (D-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LANCL3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-247406 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LANCL3 (D-17) is recommended for detection of LANCL3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with LANCL1 or LANCL2.

LANCL3 (D-17) is also recommended for detection of LANCL3 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for LANCL3 siRNA (h): sc-91362, LANCL3 siRNA (m): sc-146643, LANCL3 shRNA Plasmid (h): sc-91362-SH, LANCL3 shRNA Plasmid (m): sc-146643-SH, LANCL3 shRNA (h) Lentiviral Particles: sc-91362-V and LANCL3 shRNA (m) Lentiviral Particles: sc-146643-V.

Molecular Weight of LANCL3 isoforms: 46/43 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.