

LOC100130172 (G-20): sc-247494

BACKGROUND

LOC100130172 is a 173 amino acid protein that is encoded by a gene that maps to human chromosome 5. Chromosome 5 contains 181 million base pairs and comprises nearly 6% of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm or of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

1. Edwards, S.J., et al. 1997. The mutational spectrum in Treacher Collins syndrome reveals a predominance of mutations that create a premature-termination codon. *Am. J. Hum. Genet.* 60: 515-524.
2. McDaniel, L.D., et al. 1997. Confirmation of homozygosity for a single nucleotide substitution mutation in a Cockayne syndrome patient using monoallelic mutation analysis in somatic cell hybrids. *Hum. Mutat.* 10: 317-321.
3. Crawford, M.J., et al. 1997. Human and murine PTX1/Ptx1 gene maps to the region for Treacher Collins syndrome. *Mamm. Genome* 8: 841-845.
4. Finch, R., et al. 2005. Familial adenomatous polyposis and mental retardation caused by a *de novo* chromosomal deletion at 5q15-q22: report of a case. *Dis. Colon Rectum.* 48: 2148-2152.
5. Anindya, R., et al. 2007. Damage-induced ubiquitylation of human RNA polymerase II by the ubiquitin ligase Nedd4, but not Cockayne syndrome proteins or BRCA1. *Mol. Cell* 28: 386-397.
6. Vera-Carbonell, A., et al. 2009. Characterization of a *de novo* complex chromosomal rearrangement in a patient with cri-du-chat and trisomy 5p syndromes. *Am. J. Med. Genet. A* 149A: 2513-2521.
7. Ravandi, F., et al. 2009. Superior outcome with hypomethylating therapy in patients with acute myeloid leukemia and high-risk myelodysplastic syndrome and chromosome 5 and 7 abnormalities. *Cancer* 115: 5746-5751.
8. Sazawal, S., et al. 2009. Haematological & molecular profile of acute myelogenous leukaemia in India. *Indian J. Med. Res.* 129: 256-261.

CHROMOSOMAL LOCATION

Genetic locus: LOC100130172 (human) mapping to 5q31.2.

SOURCE

LOC100130172 (G-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LOC100130172 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-247494 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

LOC100130172 (G-20) is recommended for detection of LOC100130172 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of LOC100130172: 22 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.