

SATL1 (A-20): sc-248534

BACKGROUND

SATL1 (spermine N(1)-acetyltransferase-like protein 1) is a 508 amino acid protein that contains one N-acetyltransferase domain, and belongs to the acetyltransferase family. Existing as two alternatively spliced isoforms, the SATL1 gene is conserved in chimpanzee, canine, bovine, mouse and rat, and maps to human chromosome Xq21.1. Chromosome X consists of about 153 million base pairs and nearly 1,000 genes. The combination of an X and Y chromosome lead to normal male development while two copies of X lead to normal female development. There are a number of conditions related to an unusual number and combination of sex chromosomes being inherited. More than one copy of the X chromosome with a Y chromosome causes Klinefelter's syndrome. A single copy of X alone leads to Turner's syndrome. More than two copies of the X chromosome, in the absence of a Y chromosome, is known as triple X syndrome. Color blindness, hemophilia and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently as males carry a single X chromosome.

REFERENCES

- Bernardino-Sgherri, et al. 2002. Overall DNA methylation and chromatin structure of normal and abnormal X chromosomes. *Cytogenet. Genome Res.* 99: 85-91.
- Deeb, S.S. 2005. The molecular basis of variation in human color vision. *Clin. Genet.* 67: 369-377.
- Ross, M.T., et al. 2005. The DNA sequence of the human X chromosome. *Nature* 434: 325-337.
- Hayashi, T., et al. 2006. Novel form of a single X-linked visual pigment gene in a unique dichromatic color-vision defect. *Vis. Neurosci.* 23: 411-417.
- Augui, S., et al. 2007. Sensing X chromosome pairs before X inactivation via a novel X-pairing region of the Xic. *Science* 318: 1632-1636.
- Yaspan, B.L., et al. 2008. A haplotype at chromosome Xq27.2 confers susceptibility to prostate cancer. *Hum. Genet.* 123: 379-386.
- Bahi-Buisson, N., et al. 2010. Epileptic encephalopathy in a girl with an interstitial deletion of Xp22 comprising promoter and exon 1 of the CDKL5 gene. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 153B: 202-207.

CHROMOSOMAL LOCATION

Genetic locus: SATL1 (human) mapping to Xq21.1.

SOURCE

SATL1 (A-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of SATL1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-248534 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SATL1 (A-20) is recommended for detection of SATL1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with SAT-1, SATB and SATB2.

Suitable for use as control antibody for SATL1 siRNA (h): sc-106534, SATL1 shRNA Plasmid (h): sc-106534-SH and SATL1 shRNA (h) Lentiviral Particles: sc-106534-V.


Molecular Weight of SATL1 isoforms 1/2: 56/49 kDa.

Positive Controls: A-431 whole cell lysate: sc-22201.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SATL1 (A-20): sc-248534. Western blot analysis of SATL1 expression in A-431 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **SATL1 (E-8): sc-398793**, our highly recommended monoclonal alternative to SATL1 (A-20).