

TSPAN11 (S-17): sc-249109

BACKGROUND

Tetraspanins are a group of hydrophobic membrane proteins that interact with a wide variety of proteins including intracellular signaling molecules, integrins and membrane receptors. Members of the tetraspanin family are characterized by the presence of four hydrophobic domains and play a role in cell development, activation, growth and motility. TSPAN11 (tetraspanin-11) is a 253 amino acid multi-pass membrane protein belonging to the tetraspanin (TM4SF) family. The gene encoding TSPAN11 maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondro-genesis, Kniest dysplasia, Noonan syndrome and trisomy 12p, which causes facial developmental defects and seizure disorders.

REFERENCES

- Todd, S.C., et al. 1998. Sequences and expression of six new members of the tetraspanin/TM4SF family. *Biochim. Biophys. Acta* 1399: 101-104.
- Domínguez-Jimenez, C., et al. 2001. Involvement of $\alpha 3$ integrin/tetraspanin complexes in the angiogenic response induced by angiotensin II. *FASEB J.* 15: 1457-1459.
- Berditchevski, F. 2001. Complexes of tetraspanins with integrins: more than meets the eye. *J. Cell Sci.* 114: 4143-4151.
- Yokoyama, T., et al. 2003. A case of kniest dysplasia with retinal detachment and the mutation analysis. *Am. J. Ophthalmol.* 136: 1186-1188.
- Chen, L., et al. 2008. Clinicopathological significance of overexpression of TSPAN1, Ki67 and CD34 in gastric carcinoma. *Tumori* 94: 531-538.
- Benussi, D.G., et al. 2009. Trisomy 12p and monosomy 4p: phenotype-genotype correlation. *Genet. Test. Mol. Biomarkers* 13: 199-204.
- Scholz, C.J., et al. 2009. Tspan-1 is a tetraspanin preferentially expressed by mucinous and endometrioid subtypes of human ovarian carcinomas. *Cancer Lett.* 275: 198-203.
- Chen, L., et al. 2009. TSPAN1 protein expression: a significant prognostic indicator for patients with colorectal adenocarcinoma. *World J. Gastroenterol.* 15: 2270-2276.
- Bennett, G., et al. 2011. A functional and transcriptomic analysis of NET1 bioactivity in gastric cancer. *BMC Cancer* 11: 50.

CHROMOSOMAL LOCATION

Genetic locus: Tspan11 (mouse) mapping to 6 F3.

SOURCE

TSPAN11 (S-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TSPAN11 of mouse origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-249109 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TSPAN11 (S-17) is recommended for detection of TSPAN11 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other TSPAN family members.

TSPAN11 (S-17) is also recommended for detection of TSPAN11 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for TSPAN11 siRNA (m): sc-154724, TSPAN11 shRNA Plasmid (m): sc-154724-SH and TSPAN11 shRNA (m) Lentiviral Particles: sc-154724-V.

Molecular Weight of TSPAN11: 28 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.