

UGT3A2 (T-16): sc-249225

BACKGROUND

UDP-glucuronosyltransferase isoenzymes (UGTs) catalyze the glucuronidation of small lipophilic molecules, thereby regulating the bioactivity and metabolic fate of a wide range of endogenous compounds and xenobiotics. Glucuronidation increases the polarity of lipophilic molecules and facilitates their entry into aqueous compartments and, ultimately, their excretion. In essence, glucuronidation provides a protective function by terminating or attenuating the biological activity of its substrates. The UGT3A family of isoenzymes are expressed in liver and kidney, and to a lesser, in the gastrointestinal tract. UGT3A2 (UDP glycosyltransferase 3 family, polypeptide A2) is a 523 amino acid single-pass type I membrane protein that belongs to the UDP-glycosyltransferase family. It has been suggested that members of the UGT3A family may have an important role in the metabolism and elimination of ursodeoxycholic acid, a metabolic byproduct of intestinal bacteria.

REFERENCES

1. Bélanger, A., Hum, D.W., Beaulieu, M., Lévesque, E., Guillemette, C., Tcherno, A., Bélanger, G., Turgeon, D. and Dubois, S. 1998. Characterization and regulation of UDP-glucuronosyltransferases in steroid target tissues. *J. Steroid Biochem. Mol. Biol.* 65: 301-310.
2. Lévesque, E., Beaulieu, M., Hum, D.W. and Bélanger, A. 1999. Characterization and substrate specificity of UGT2B4 (E458): a UDP-glucuronosyltransferase encoded by a polymorphic gene. *Pharmacogenetics* 9: 207-216.
3. King, C.D., Rios, G.R., Green, M.D. and Tephly, T.R. 2000. UDP-glucuronosyltransferases. *Curr. Drug Metab.* 1: 143-161.
4. Mackenzie, P.I., Bock, K.W., Burchell, B., Guillemette, C., Ikushiro, S., Iyanagi, T., Miners, J.O., Owens, I.S. and Nebert, D.W. 2005. Nomenclature update for the mammalian UDP glycosyltransferase (UGT) gene superfamily. *Pharmacogenet. Genomics* 15: 677-685.
5. Buckley, D.B. and Klaassen, C.D. 2007. Tissue- and gender-specific mRNA expression of UDP-glucuronosyltransferases (UGTs) in mice. *Drug Metab. Dispos.* 35: 121-127.
6. Argikar, U.A., Iwuchukwu, O.F. and Nagar, S. 2008. Update on tools for evaluation of uridine diphosphoglucuronosyltransferase polymorphisms. *Expert Opin. Drug Metab. Toxicol.* 4: 879-894.
7. Mackenzie, P.I., Rogers, A., Treloar, J., Jorgensen, B.R., Miners, J.O. and Meech, R. 2008. Identification of UDP glycosyltransferase 3A1 as a UDP N-acetylglucosaminyltransferase. *J. Biol. Chem.* 283: 36205-36210.

CHROMOSOMAL LOCATION

Genetic locus: UGT3A2 (human) mapping to 5p13.2.

SOURCE

UGT3A2 (T-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of UGT3A2 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-249225 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

UGT3A2 (T-16) is recommended for detection of UGT3A2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with UGT3A1.

Suitable for use as control antibody for UGT3A2 siRNA (h): sc-91750, UGT3A2 shRNA Plasmid (h): sc-91750-SH and UGT3A2 shRNA (h) Lentiviral Particles: sc-91750-V.

Molecular Weight of UGT3A2: 60 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.