

ZF5128 (L-17): sc-249415

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZF5128, also known as zinc finger protein 324A or zinc finger protein ZF5128, is a 553 amino acid protein that localizes to the nucleus and is composed of nine C₂H₂-type zinc fingers and one KRAB domain. ZF5128 is suggested to play a role in T cell activation. ZF5128 is encoded by a gene located on human chromosome 19, which consists of around 63 million bases with over 1,400 genes and makes up over 2% of human genomic DNA.

REFERENCES

1. Rousseau-Merck, M.F., et al. 1993. Chromosomal localization of 9 KOX zinc finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. *Hum. Genet.* 92: 583-587.
2. Urrutia, R. 2003. KRAB-containing zinc-finger repressor proteins. *Genome Biol.* 4: 231.
3. Sun, Y., et al. 2003. The KRAB domain of zinc finger gene ZNF268: a potential transcriptional repressor. *IUBMB Life* 55: 127-131.
4. Englbrecht, C.C., et al. 2004. Conservation, diversification and expansion of C₂H₂ zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39.
5. Nakamura, M., et al. 2004. A novel subfamily of zinc finger genes involved in embryonic development. *J. Cell. Biochem.* 93: 887-895.
6. Zhong, Z., et al. 2007. Identification of a novel human zinc finger gene, ZNF438, with transcription inhibition activity. *J. Biochem. Mol. Biol.* 40: 517-524.
7. O'Geen, H., et al. 2007. Genome-wide analysis of KAP1 binding suggests autoregulation of KRAB-ZNFs. *PLoS Genet.* 3: e89.

CHROMOSOMAL LOCATION

Genetic locus: Zfp324 (mouse) mapping to 7 A1.

SOURCE

ZF5128 (L-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZF5128 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-249415 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ZF5128 (L-17) is recommended for detection of ZF5128 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZNF family members.

Suitable for use as control antibody for ZF5128 siRNA (m): sc-155510, ZF5128 shRNA Plasmid (m): sc-155510-SH and ZF5128 shRNA (m) Lentiviral Particles: sc-155510-V.

Molecular Weight of ZF5128: 61 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.