

ZNF10 (P-17): sc-249447

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF10 (zinc finger protein 10), also known as KOX1, is a 573 amino acid protein that localizes to the nucleus and contains one KRAB domain and 11 CHH2-type zinc fingers. One of several members of the Krüppel C₂H₂-type zinc-finger protein family, ZNF10 exists as multiple alternatively spliced isoforms and is thought to play a role in transcriptional regulation. The gene encoding ZNF10 maps to human chromosome 12, which houses over 1,100 genes and comprises approximately 4.5% of the human genome.

REFERENCES

1. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
2. Huebner, K., Druck, T., Croce, C.M. and Thiesen, H.J. 1991. Twenty-seven nonoverlapping zinc finger cDNAs from human T cells map to nine different chromosomes with apparent clustering. *Am. J. Hum. Genet.* 48: 726-740.
3. Rousseau-Merck, M.F., Hillion, J., Jonveaux, P., Couillin, P., Seite, P., Thiesen, H.J. and Berger, R. 1993. Chromosomal localization of 9 KOX zinc finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. *Hum. Genet.* 92: 583-587.
4. Moosmann, P., Georgiev, O., Le Douarin, B., Bourquin, J.P. and Schaffner, W. 1996. Transcriptional repression by RING finger protein TIF1 β that interacts with the KRAB repressor domain of KOX1. *Nucleic Acids Res.* 24: 4859-4867.
5. Moosmann, P., Georgiev, O., Thiesen, H.J., Hagmann, M. and Schaffner, W. 1997. Silencing of RNA polymerases II and III-dependent transcription by the KRAB protein domain of KOX1, a Krüppel-type zinc finger factor. *Biol. Chem.* 378: 669-677.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 194538. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: ZNF10 (human) mapping to 12q24.33.

SOURCE

ZNF10 (P-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZNF10 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-249447 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ZNF10 (P-17) is recommended for detection of ZNF10 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ZNF10 (P-17) is also recommended for detection of ZNF10 in additional species, including bovine.

Suitable for use as control antibody for ZNF10 siRNA (h): sc-95814, ZNF10 shRNA Plasmid (h): sc-95814-SH and ZNF10 shRNA (h) Lentiviral Particles: sc-95814-V.

Molecular Weight of ZNF10: 50 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.