

Dicer (C-20): sc-25117

BACKGROUND

The mammalian Dicer is a type III RNase-related protein with orthologs in yeast, *Drosophila* and *Arabidopsis*. Dicer contains an RNA-helicase motif, including a DEXH box in its amino-terminus and an RNase motif in the carboxy-terminus. The gene encoding human Dicer maps to chromosome 14q32.13. Dicer is expressed in brain, heart, liver, lung, pancreas, kidney and placenta, and functions in the RNA interference pathway. Dicer cleaves short hairpin RNA precursors of approximately 70 bp into 21-23 bp dsRNAs that selectively target the destruction of homologous RNAs. Dicer localizes to the cytoplasm of mammalian cells. Specifically, it co-localizes with calreticulin in the endoplasmic reticulum. Although the cleavage of RNA by Dicer is ATP-independent, the product release necessary for the rapid turnover of this enzyme may be attributed to ATP. Immunoprecipitation studies indicate Dicer forms a complex with the PIWI domain of eIF2C translation initiation factors.

CHROMOSOMAL LOCATION

Genetic locus: DICER1 (human) mapping to 14q32.13; Dicer1 (mouse) mapping to 12 E.

SOURCE

Dicer (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Dicer of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-25117 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as agarose conjugate for immunoprecipitation, sc-25117 AC, 500 µg/0.25 ml agarose in 1 ml.

APPLICATIONS

Dicer (C-20) is recommended for detection of Dicer of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Dicer (C-20) is also recommended for detection of Dicer in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Dicer siRNA (h): sc-40489, Dicer siRNA (m): sc-40490, Dicer shRNA Plasmid (h): sc-40489-SH, Dicer shRNA Plasmid (m): sc-40490-SH, Dicer shRNA (h) Lentiviral Particles: sc-40489-V and Dicer shRNA (m) Lentiviral Particles: sc-40490-V.


Molecular Weight of Dicer: 218 kDa.

Positive Controls: rat brain extract: sc-2392, HeLa whole cell lysate: sc-2200 or Hep G2 cell lysate: sc-2227.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Dicer (C-20): sc-25117. Western blot analysis of Dicer expression in rat brain tissue extract.



Dicer (C-20): sc-25117. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

- Zhao, Y., et al. 2007. Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking miRNA-1-2. *Cell* 129: 303-317.
- Nishi, H., et al. 2011. MicroRNA-27a regulates β cardiac myosin heavy chain gene expression by targeting thyroid hormone receptor β 1 in neonatal rat ventricular myocytes. *Mol. Cell. Biol.* 31: 744-755.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Dicer (F-10): sc-136979** or **Dicer (E-7): sc-393328**, our highly recommended monoclonal alternatives to Dicer (C-20). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Dicer (F-10): sc-136979**.