# SANTA CRUZ BIOTECHNOLOGY, INC.

# Bag-1 (F-7): sc-25296



#### BACKGROUND

The Bcl-2 family of proteins is characterized by its ability to modulate cell death (apoptosis) under a broad range of physiologic conditions. Bcl-2 and several related proteins function to inhibit apoptosis, while other members of the Bcl-2 family, such as Bax and Bak, enhance cell death under various conditions. For instance, Bcl-x<sub>L</sub> represses cell death, while its shorter form, Bcl-x<sub>S</sub>, promotes apoptosis. Dimerization of another member of this family, Bad, with Bcl-x<sub>L</sub> results in displacement of Bax from Bcl-x<sub>L</sub>:Bax complexes and restoration of Bax-mediated apoptosis. A Bcl-2 or with other Bcl-2-related proteins. Bag-1 appears to function to enhance Bcl-2 protection from cell death, suggesting that Bag-1 represents a new type of anti-cell death gene and that certain routes of apoptosis induction previously ascribed to Bcl-2-independent pathways may instead reflect a requirement for a combination of Bcl-2 and Bag-1.

# REFERENCES

- Nunez, G., et al. 1990. Deregulated Bcl-2 gene expression selectively prolongs survival of growth factor-deprived hemopoietic cell lines. J. Immunol. 144: 3602-3610.
- Hockenbery, D.M., et al. 1991. Bcl-2 protein is topographically restricted in tissues characterized by apoptotic cell death. Proc. Natl. Acad. Sci. USA 88: 6961-6965.

## CHROMOSOMAL LOCATION

Genetic locus: BAG1 (human) mapping to 9p13.3.

#### SOURCE

Bag-1 (F-7) is a mouse monoclonal antibody raised against amino acids 73-274 of Bag-1 of human origin.

## PRODUCT

Each vial contains 200  $\mu g$  IgG\_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

#### **APPLICATIONS**

Bag-1 (F-7) is recommended for detection of Bag-1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:500), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Bag-1 siRNA (h): sc-29211, Bag-1 shRNA Plasmid (h): sc-29211-SH and Bag-1 shRNA (h) Lentiviral Particles: sc-29211-V.

Molecular Weight of four major isoforms of Bag-1: 32/36/46/50 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or Bag-1 (h): 293T Lysate: sc-112723.

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

#### DATA





Bag-1 (F-7): sc-25296. Western blot analysis of Bag-1 expression in non-transfected: sc-117752 (**A**) and human Bag-1 transfected: sc-112723 (**B**) 293T whole cell lysates.

Bag-1 (F-7): sc-25296. Immunoperoxidase staining of formalin fixed, paraffin-embedded human breast tissue showing cytoplasmic and nuclear staining of qlandular cells.

#### SELECT PRODUCT CITATIONS

- Maddalo, D., et al. 2012. A peptidic unconjugated GRP78/BiP ligand modulates the unfolded protein response and induces prostate cancer cell death. PLoS ONE 7: e45690.
- 2. Cato, L., et al. 2017. Development of Bag-1L as a therapeutic target in androgen receptor-dependent prostate cancer. Elife 6: e27159.

#### **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.