

## IL-10R $\alpha$ (H-300): sc-25477

### BACKGROUND

The IL-10 receptor, IL-10R, is a member of the class II subgroup of the cytokine receptor family and exhibits structural similarity to the interferon receptor. IL-10R is expressed in B cells and T helper cells, as well as in LPS-induced mouse fibroblasts. Overall, mouse IL-10R and human IL-10R share 60% sequence identity at the protein level. Stimulation with IL-10 leads to phosphorylation of JAK1 and Tyk 2 tyrosine kinases. The activated kinases phosphorylate the two tyrosine residues (Tyr 446 and Tyr 496) in the cytoplasmic domain of IL-10R $\alpha$ . The phosphorylation of these two residues are required for proper function of IL-10R and activation of IL-10E1 signaling. IL-10R $\beta$  is ubiquitously expressed and, in addition to forming the IL-10 heterodimeric receptor, it forms a heterodimeric receptor with an IL-22R subunit and an IL-28R subunit. IL-10R is constitutively expressed on human natural killer (NK) cells and the direct binding of IL-10 potentiates cytokine production by human NK cells.

### REFERENCES

1. Rousset, F., Garcia, E., Defrance, T., Péronne, C., Vezzio, N., Hsu, D.-H., Kastelein, R., Moore, K.W. and Banchereau, J. 1992. Interleukin 10 is a potent growth and differentiation factor for activated human B lymphocytes. *Proc. Natl. Acad. Sci. USA* 89: 1890-1893.
2. Ho, A.S.Y., Liu, Y., Khan, T.A., Hsu, D.-H., Bazan, J.F. and Moore, K.W. 1993. A receptor for interleukin 10 is related to interferon receptors. *Proc. Natl. Acad. Sci. USA* 90: 11267-11271.
3. Liu, Y., Wei, S.H., Ho, A.S., de Waal Malefyt, R. and Moore, K.W. 1994. Expression cloning and characterization of a human IL-10 receptor. *J. Immunol.* 152: 1821-1829.
4. Weber-Nordt, R.M., Meraz, M.A. and Schreiber, R.D. 1994. Lipopolysaccharide-dependent induction of IL-10 receptor expression on murine fibroblasts. *J. Immunol.* 153: 3734-3744.
5. Ho, A.S., Wei, S.H., Mui, A.L., Miyajima, A. and Moore, K.W. 1995. Functional regions of the mouse interleukin-10 receptor cytoplasmic domain. *Mol. Cell. Biol.* 15: 5043-5053.
6. Tan, J.C., Braun, S., Rong, H., DiGiacomo, R., Dolphin, E., Baldwin, S., Narula, S.K., Zavodny, P.J. and Chou, C.C. 1995. Characterization of recombinant extracellular domain of human interleukin-10 receptor. *J. Biol. Chem.* 270: 12906-12911.
7. Carson, W.E., Lindemann, M.J., Baiocchi, R., Linett, M., Tan, J.C., Chou, C.C., Narula, S. and Caligiuri, M.A. 1995. The functional characterization of interleukin-10 receptor expression on human natural killer cells. *Blood* 85: 3577-3585.

### SOURCE

IL-10R $\alpha$  (H-300) is a rabbit polyclonal antibody raised against amino acids 1-300 of IL-10R $\alpha$  of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

IL-10R $\alpha$  (H-300) is recommended for detection of IL-10R $\alpha$  of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-10R $\alpha$  siRNA (h): sc-35646, IL-10R $\alpha$  shRNA Plasmid (h): sc-35646-SH and IL-10R $\alpha$  shRNA (h) Lentiviral Particles: sc-35646-V.

Molecular Weight of IL-10R $\alpha$ : 90-110 kDa.

Positive Controls: Ramos cell lysate: sc-2216, NAMALWA cell lysate: sc-2234 or CCRF-HSB-2 cell lysate: sc-2265.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **IL-10R $\alpha$  (A-3): sc-365374** or **IL-10R $\alpha$  (E-6): sc-28371**, our highly recommended monoclonal alternatives to IL-10R $\alpha$  (H-300).