

# PC-PLD2 (H-133): sc-25513

## BACKGROUND

Virtually every cell uses phosphatidylcholine as a substrate to produce phosphatidic acid and choline. Phosphatidylcholine phospholipase D1 and D2 (PC-PLD1 and PC-PLD2) are phospholipid-specific phosphodiesterases that hydrolyze phosphatidylcholine. Unlike PC-PLD1, which associates with secretory granules, PC-PLD2 localizes to the plasma membrane, where it is implicated in the formation of endocytotic vesicles. Both PC-PLD1 and PC-PLD2 coordinately regulate macrophage phagocytosis. PC-PLD activity in mammalian cells is transiently stimulated upon activation by G protein-coupled and receptor tyrosine kinase cell surface receptors. For example, PC-PLD1 and PC-PLD2 participate in sphingosine 1-phosphate stimulation of ERK phosphorylation and IL-8 secretion in bronchial epithelial cells. In addition, Tubulin binding to PC-PLD2 inhibits muscarinic receptor-linked PC-PLD2 activation. PC-PLD2 also enhances PKC $\zeta$  activity through direct interaction in a lipase activity-independent manner. PC-PLD1 and PC-PLD2 stimulate cell growth by repressing expression of p21 gene through p53-dependent and p53-independent pathways, respectively, which may ultimately lead to carcinogenesis.

## CHROMOSOMAL LOCATION

Genetic locus: PLD2 (human) mapping to 17p13.2; Pld2 (mouse) mapping to 11 B3.

## SOURCE

PC-PLD2 (H-133) is a rabbit polyclonal antibody raised against amino acids 8-140 of PC-PLD2 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

PC-PLD2 (H-133) is recommended for detection of PC-PLD2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PC-PLD2 (H-133) is also recommended for detection of PC-PLD2 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for PC-PLD2 siRNA (h): sc-44001, PC-PLD2 siRNA (m): sc-61367, PC-PLD2 shRNA Plasmid (h): sc-44001-SH, PC-PLD2 shRNA Plasmid (m): sc-61367-SH, PC-PLD2 shRNA (h) Lentiviral Particles: sc-44001-V and PC-PLD2 shRNA (m) Lentiviral Particles: sc-61367-V.


Molecular Weight of PC-PLD2: 117 kDa.

Positive Controls: U-937 cell lysate: sc-2239.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



PC-PLD2 (H-133): sc-25513. Western blot analysis of PC-PLD2 expression in U-937 whole cell lysate.

## SELECT PRODUCT CITATIONS

- Disse, J., et al. 2009. Phospholipase D1 is specifically required for regulated secretion of von Willebrand factor from endothelial cells. *Blood* 113: 973-980.
- Krishnan, B., et al. 2011. Dopamine-induced plasticity, phospholipase D (PLD) activity and cocaine-cue behavior depend on PLD-linked metabotropic glutamate receptors in amygdala. *PLoS ONE* 6: e25639.
- Chen, F., et al. 2013. Phospholipase D2 mediates signaling by ATPase class I type 8B membrane 1. *J. Lipid Res.* 54: 379-385.
- Li, S., et al. 2013. The neural cell adhesion molecule (NCAM) associates with and signals through p21-activated kinase 1 (Pak1). *J. Neurosci.* 33: 790-803.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **PC-PLD2 (1C5): sc-293214**, our highly recommended monoclonal alternative to PC-PLD2 (H-133).