

PDE7A1/2 (H-52): sc-25566

BACKGROUND

Phosphodiesterases (PDE, also designated cyclic nucleotide phosphodiesterase) are important for the downregulation of the intracellular level of the second messenger cyclic adenosine monophosphate (cAMP) by hydrolyzing cAMP to 5'AMP. Phosphodiesterase type 3 isoforms, PDE3A and 3B, are expressed primarily in cardiovascular tissue and adipose tissue, respectively. PDE3A, is found in myocardium and platelets and PDE3B is found in lymphocytes. The PDE7A1 (HCP1) isozyme and the PDE7A2 proteins, alternate splice products of PDE7A, are highly expressed in skeletal muscle. PDE7B is most highly expressed in pancreas. The PDE family contains proteins that serve tissue-specific roles in regulation of lipolysis, glycogenolysis, myocardial contractility, and smooth muscle relaxation.

REFERENCES

1. Bloom, T.J. and Beavo, J.A. 1996. Identification and tissue-specific expression of PDE7 phosphodiesterase splice variants. *Proc. Natl. Acad. Sci. USA* 93: 14188-14192.
2. Han, P., et al. 1997. Alternative splicing of the high affinity cAMP-specific phosphodiesterase (PDE7A) mRNA in human skeletal muscle and heart. *J. Biol. Chem.* 272: 16152-16157.

CHROMOSOMAL LOCATION

Genetic locus: PDE7A (human) mapping to 8q13.1; Pde7a (mouse) mapping to 3 A2.

SOURCE

PDE7A1/2 (H-52) is a rabbit polyclonal antibody raised against amino acids 431-482 of PDE7A1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PDE7A1/2 (H-52) is recommended for detection of PDE7A1 and PDE7A2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PDE7A1/2 (H-52) is also recommended for detection of PDE7A1 and PDE7A2 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for PDE7A siRNA (h): sc-44005, PDE7A siRNA (m): sc-41609, PDE7A shRNA Plasmid (h): sc-44005-SH, PDE7A shRNA Plasmid (m): sc-41609-SH, PDE7A shRNA (h) Lentiviral Particles: sc-44005-V and PDE7A shRNA (m) Lentiviral Particles: sc-41609-V.


Molecular Weight of PDE7A1/2: 57/50 kDa.

Positive Controls: HuT 78 whole cell lysate: sc-2208.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PDE7A1/2 (H-52): sc-25566. Western blot analysis of PDE7A1/2 expression in HuT 78 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Smith, S.J., et al. 2003. Ubiquitous expression of phosphodiesterase 7A in human proinflammatory and immune cells. *Am. J. Physiol.* 284: L279-L289.
2. Kadoshima-Yamaoka, K., et al. 2009. ASB16165, a phosphodiesterase 7A inhibitor, reduces cutaneous TNF-α level and ameliorates skin edema in phorbol ester 12-O-tetradecanoylphorbol-13-acetate-induced skin inflammation model in mice. *Eur. J. Pharmacol.* 613: 163-166.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **PDE7A (B-11): sc-398031** or **PDE7A1/2 (H-7): sc-28374**, our highly recommended monoclonal alternatives to PDE7A1/2 (H-52).