

# SLAM (H-80): sc-25611

## BACKGROUND

Following occupancy of the T cell receptor by antigen, T cell proliferation and lymphokine production are determined by a second costimulatory signal delivered by a ligand expressed on antigen-presenting cells. SLAM (for signaling lymphocyte-activation molecule, also designated CDw150) is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. SAP (for SLAM-associated protein) contains an SH2 domain and functions to inhibit SH-PTP2 recruitment to the SLAM docking site, an activity induced by Fyn phosphorylation of SLAM. Mutations of the SAP gene may be associated with X-linked lymphoproliferative disease (XLP).

## REFERENCES

- Freeman, G.J., Gray, G.S., Gimmi, C.D., Lombard, D.B., Zhou, L.J., White, M., Fingerth, J.D., Gribben, J.G. and Nadler, L.M. 1991. Structure, expression, and T cell costimulatory activity of the murine homologue of the human B lymphocyte activation antigen B7. *J. Exp. Med.* 174: 625-631.
- Cocks, B.G., Chang, C.C., Carballido, J.M., Yssel, H., de Vries, J.E. and Aversa, G. 1995. A novel receptor involved in T cell activation. *Nature* 376: 260-263.
- Aversa, G., Carballido, J., Punnonen, J., Chang, C.C., Hauser, T., Cocks, B.G. and De Vries, J.E. 1997. SLAM and its role in T cell activation and T cell responses. *Immunol. Cell Biol.* 75: 202-205.
- Aversa, G., Chang, C.C., Carballido, J.M., Cocks, B.G. and De Vries, J.E. 1997. Engagement of the signaling lymphocytic activation molecule (SLAM) on activated T cells results in IL-2-independent, cyclosporin A-sensitive T cell proliferation and IFN- $\gamma$  production. *J. Immunol.* 158: 4036-4044.
- Favero, J. and Lafont, V. 1998. Effector pathways regulating T cell activation. *Biochem. Pharmacol.* 56: 1539-1547.
- Sayos, J., Wu, C., Morra, M., Wang, N., Zhang, X., Allen, D., van Schaik, S., Notarangelo, L., Geha, R., Roncarolo, M.G., Oettgen, H., De Vries, J.E., Aversa, G. and Terhorst, C. 1998. The X-linked lymphoproliferative-disease gene product SAP regulates signals induced through the co-receptor SLAM. *Nature* 395: 462-469.

## CHROMOSOMAL LOCATION

Genetic locus: SLAMF1 (human) mapping to 1q23.3.

## SOURCE

SLAM (H-80) is a rabbit polyclonal antibody raised against amino acids 31-110 mapping near the N-terminus of SLAM of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

SLAM (H-80) is recommended for detection of SLAM of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SLAM siRNA (h): sc-42974, SLAM shRNA Plasmid (h): sc-42974-SH and SLAM shRNA (h) Lentiviral Particles: sc-42974-V.


Molecular Weight of SLAM: 70 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, MOLT-4 cell lysate: sc-2233 or HuT 78 whole cell lysate: sc-2208.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



SLAM (H-80): sc-25611. Western blot analysis of SLAM expression in Jurkat whole cell lysate.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

**MONOS**  
Satisfaction  
Guaranteed

Try **SLAM (E-11): sc-166939**, our highly recommended monoclonal alternative to SLAM (H-80).