

P2X6 (H-81): sc-25697

BACKGROUND

The P2X receptor family is comprised of ligand-gated ion channels that allow for the increased permeability of calcium into the cell in response to extracellular ATP. There are seven P2X receptors, P2X1, P2X2, P2X3, P2X4, P2X5, P2X6, P2X7, which form either homomeric or heteromeric channels or both, and they are characterized by intracellular amino and carboxy termini. P2X receptors are expressed in a wide variety of tissues, including neurons, prostate, bladder, pancreas, colon, testis, and ovary. The major function of the P2X receptors is to mediate synaptic transmissions between neurons and to other tissues via the binding of extracellular ATP, which acts as a neurotransmitter. The P2X receptors may be involved in the onset of necrosis or apoptosis after prolonged exposure to high concentrations of extracellular ATP.

REFERENCES

1. Longhurst, P.A., et al. 1996. The human P2X1 receptor: molecular cloning, tissue distribution, and localization to chromosome 17. *Biochim. Biophys. Acta* 1308: 185-188.
2. Di Virgilio, F., et al. 1998. Cytolytic P2X purinoceptors. *Cell Death Differ.* 5: 191-199.
3. Alexander, K., et al. 1999. Allosteric modulation and accelerated resensitization of human P2X3 receptors by cibacron blue. *J. Pharmacol. Exp. Ther.* 291: 1135-1142.
4. Burnstock, G. 2000. P2X receptors in sensory neurones. *Br. J. Anaesth.* 84: 476-488.
5. Oury, C., et al. 2000. A natural dominant negative P2X1 receptor due to deletion of a single amino acid residue. *J. Biol. Chem.* 275: 22611-22614.
6. Ding, S., et al. 2000. Inactivation of P2X2 purinoceptors by divalent cations. *J. Physiol.* 522: 199-214.
7. North, R.A., et al. 2000. Pharmacology of cloned P2X receptors. *Annu. Rev. Pharmacol. Toxicol.* 40: 563-580.

CHROMOSOMAL LOCATION

Genetic locus: P2RX6 (human) mapping to 22q11.21; P2rx6 (mouse) mapping to 16 A3.

SOURCE

P2X6 (H-81) is a rabbit polyclonal antibody raised against amino acids 351-431 mapping at the C-terminus of P2X6 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

P2X6 (H-81) is recommended for detection of P2X6 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for P2X6 siRNA (h): sc-42573, P2X6 siRNA (m): sc-42574, P2X6 shRNA Plasmid (h): sc-42573-SH, P2X6 shRNA Plasmid (m): sc-42574-SH, P2X6 shRNA (h) Lentiviral Particles: sc-42573-V and P2X6 shRNA (m) Lentiviral Particles: sc-42574-V.


Molecular Weight of P2X6: 49-70 kDa, depending on glycosylation.

Positive Controls: P2X6 (h): 293T Lysate: sc-112163 or L8 cell lysate: sc-3807.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



P2X6 (H-81): sc-25697. Western blot analysis of P2X6 expression in non-transfected: sc-117752 (A) and human P2X6 transfected: sc-112163 (B) 293T whole cell lysates.



P2X6 (H-81): sc-25697. Western blot analysis of P2X6 expression in L8 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Liang, L., et al. 2005. RNA interference targeted to multiple P2X receptor subtypes attenuates zinc-induced calcium entry. *Am. J. Physiol., Cell Physiol.* 289: C388-C396.

MONOS
Satisfaction
Guaranteed

Try **P2X6 (D-1): sc-166013** or **P2X6 (H-6): sc-166014**, our highly recommended monoclonal alternatives to P2X6 (H-81).