

CLC-1 (H-110): sc-25699

BACKGROUND

The family of voltage-dependent chloride channels (CLCs) regulate cellular trafficking of chloride ions, a critical component of all living cells. CLCs regulate excitability in muscle and nerve cells, aid in organic solute transport and maintain cellular volume. CLC1 is highly expressed in skeletal muscle. Mutations in the gene encoding CLC1 lead to myotonia, an inheritable disorder characterized by muscle stiffness and renal salt wasting. CLC2 is highly expressed in the epithelia of several organs including lung, which suggests CLC2 may be a possible therapeutic target for cystic fibrosis. CLC3 expression is particularly abundant in neuronal tissue, while CLC4 expression is evident in skeletal and cardiac muscle as well as brain. Mutations in the gene encoding CLC5 lead to Dent's disease, a renal disorder characterized by proteinuria and hypercalciuria. CLC6 and CLC7 are broadly expressed in several tissues including testis, kidney, brain and muscle.

REFERENCES

- Koch, M.C., Steinmeyer, K., Lorenz, C., Ricker, K., Wolf, F., Otto, M., Zoll, B., Lehmann-Horn, F., Grzeschik, K.H. and Jentsch, T.J. 1992. The skeletal muscle chloride channel in dominant and recessive human myotonia. *Science* 257: 797-800.
- Pook, M.A., Wrong, O., Wooding, C., Norden, A.G., Feest, T.G. and Thakker, R.V. 1993. Dent's disease, a renal Fanconi syndrome with nephrocalcinosis and kidney stones, is associated with a microdeletion involving DXS255 and maps to Xp11.22. *Hum. Mol. Genet.* 2: 2129-2134.
- van Slegtenhorst, M.A., et al. 1994. A gene from the Xp22.3 region shares homology with voltage-gated chloride channels. *Hum. Mol. Genet.* 3: 547-552.
- Borsani, G., Rugari, E.I., Tagliatela, M., Wong, C. and Ballabio, A. 1995. Characterization of a human and murine gene (CLCN3) sharing similarities to voltage-gated chloride channels and to a yeast integral membrane protein. *Genomics* 27: 131-141.
- Brandt, S. and Jentsch, T.J. 1995. CLC-6 and CLC-7 are two novel broadly expressed members of the CLC chloride channel family. *FEBS Lett.* 377: 15-20.
- Cid, L.P., Montrose-Rafizadeh, C., Smith, D.I., Guggino, W.B. and Cutting, G.R. 1995. Cloning of a putative human voltage-gated chloride channel (CLC-2) cDNA widely expressed in human tissues. *Hum. Mol. Genet.* 4: 407-413.
- Gyomai, K., Yeger, H., Ackerley, C., Garami, E. and Bear, C.E. 2000. Expression of the chloride channel CLC-2 in the murine small intestine epithelium. *Am. J. Physiol., Cell Physiol.* 279: 1787-1794.

CHROMOSOMAL LOCATION

Genetic locus: CLCN1 (human) mapping to 7q34; Clcn1 (mouse) mapping to 6 B2.1.

SOURCE

CLC-1 (H-110) is a rabbit polyclonal antibody raised against amino acids 1-110 mapping at the N-terminus of CLC-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CLC-1 (H-110) is recommended for detection of CLC-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CLC-1 siRNA (h): sc-42377, CLC-1 siRNA (m): sc-42378, CLC-1 shRNA Plasmid (h): sc-42377-SH, CLC-1 shRNA Plasmid (m): sc-42378-SH, CLC-1 shRNA (h) Lentiviral Particles: sc-42377-V and CLC-1 shRNA (m) Lentiviral Particles: sc-42378-V.


Molecular Weight of CLC-1: 124 kDa.

Positive Controls: rat brain extract: sc-2392 or A-673 cell lysate: sc-2414.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CLC-1 (H-110): sc-25699. Western blot analysis of CLC-1 expression in rat brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.