# Myosin Va (H-90): sc-25726



The Power to Question

#### **BACKGROUND**

Myosin Va, a member of the unconventional Myosin family, is a non-muscle Myosin that structurally combines elements from both nonmuscle Myosin type I and nonmuscle Myosin type II. Class V unconventional Myosins, which include Myosin Va and Myosin Vb, are nonfilamentous, Actin-binding enzymes that appear to be expressed ubiquitously. Myosin V proteins are regulated by their heavy chain phosphorylation, which occurs in the carboxy-terminal tail domain. The mouse Myosin Va gene is also known as the mouse dilute gene because mutations in this gene cause the coat color in mice to lighten. Mutations in the Myosin Va gene also result in the onset of severe neurological defects shortly after birth. Defects in the human Myosin Va gene, which maps to chromosome 15q21.2, have been implicated in Griscelli disease, a rare autosomal recessive disorder characterized by pigmentary dilution, variable cellular immunodeficiency and onset of acute, uncontrolled lymphocyte and macrophage activation.

## CHROMOSOMAL LOCATION

Genetic locus: MY05A (human) mapping to 15q21.2; Myo5a (mouse) mapping to 9 D.

#### SOURCE

Myosin Va (H-90) is a rabbit polyclonal antibody raised against amino acids 981-1070 of Myosin Va of human origin.

#### **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## **APPLICATIONS**

Myosin Va (H-90) is recommended for detection of Myosin Va of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Myosin Va (H-90) is also recommended for detection of Myosin Va in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Myosin Va siRNA (h): sc-35995, Myosin Va siRNA (m): sc-35996, Myosin Va shRNA Plasmid (h): sc-35995-SH, Myosin Va shRNA Plasmid (m): sc-35996-SH, Myosin Va shRNA (h) Lentiviral Particles: sc-35995-V and Myosin Va shRNA (m) Lentiviral Particles: sc-35996-V.


Molecular Weight of Myosin Va: 190 kDa.

Positive Controls: A-375 cell lysate: sc-3811, HeLa whole cell lysate: sc-2200 or KNRK whole cell lysate: sc-2214.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## **DATA**



Myosin Va (H-90): sc-25726. Western blot analysis of Myosin Va expression in KNRK whole cell lysate.

## **SELECT PRODUCT CITATIONS**

- Roberts, K.L. and Baines, J.D. 2010. Myosin Va enhances secretion of herpes simplex virus 1 virions and cell surface expression of viral glycoproteins. J. Virol. 84: 9889-9896.
- 2. Chaudhury, A., et al. 2011. Myosin Va plays a key role in nitrergic neurotransmission by transporting nNOS $\alpha$  to enteric varicosity membrane. Am. J. Physiol. Gastrointest. Liver Physiol. 301: G498-G507.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Myosin Va (G-4):** sc-365986, our highly recommended monoclonal alternative to Myosin Va (H-90).

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com