

PAI-2 (H-70): sc-25745

BACKGROUND

PAI-1 and PAI-2 (for plasminogen activator inhibitor-1 and -2) are members of the serpin serine proteinase inhibitor family. PAI-1 and PAI-2 have been shown to regulate uPA (urokinase-type plasminogen activator) and tPA (tissue plasminogen activator), resulting in the inhibition of proteolytic activity. Members of the serpin family generally complex with their target proteinases, then disassociate slowly into cleaved species that fold into stable inactive forms. PAI-1 can fold into the inactive state without cleavage, resulting in the latent form of PAI-1. Activity can be restored to the latent form of PAI-1 through denaturation and renaturation. PAI-2 occurs in secreted and cytosolic forms through facultative polypeptide translocation. uPA is a serine proteinase that is a member of the trypsin family. It is responsible for the cleavage of plasminogen at the Arg-Val bond to produce plasmin. uPA consists of two chains designated A and B. The A chain can be cleaved, resulting in low and high molecular mass forms of uPA.

REFERENCES

- Riccio, A., et al. 1985. The human urokinase-plasminogen activator gene and its promoter. *Nucleic Acids Res.* 13: 2759-2771.
- Belin, D., et al. 1989. Facultative polypeptide translocation allows a single mRNA to encode the secreted and cytosolic forms of plasminogen activators inhibitor 2. *EMBO J.* 8: 3287-3294.

CHROMOSOMAL LOCATION

Genetic locus: SERPINB2 (human) mapping to 18q21.33.

SOURCE

PAI-2 (H-70) is a rabbit polyclonal antibody raised against amino acids 61-130 mapping near the N-terminus of PAI-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PAI-2 (H-70) is recommended for detection of PAI-2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PAI-2 siRNA (h): sc-40804, PAI-2 shRNA Plasmid (h): sc-40804-SH and PAI-2 shRNA (h) Lentiviral Particles: sc-40804-V.

Molecular Weight of placental PAI-2: 46 kDa.


Molecular Weight of PAI-2 plasma: 60 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, CCRF-CEM cell lysate: sc-2225 or U-937 cell lysate: sc-2239.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



PAI-2 (H-70): sc-25745. Western blot analysis of PAI-2 expression in CCRF-CEM (A) and human PBL (B) whole cell lysates.



PAI-2 (H-70): sc-25745. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of exocrine glandular cells and Islets of Langerhans.

SELECT PRODUCT CITATIONS

- Boncela, J., et al. 2011. Association of plasminogen activator inhibitor type 2 (PAI-2) with proteasome within endothelial cells activated with inflammatory stimuli. *J. Biol. Chem.* 286: 43164-43171.
- Zhou, L., et al. 2013. Low expression of PAI-2 as a novel marker of portal vein tumor thrombosis and poor prognosis in hepatocellular carcinoma. *World J. Surg.* 37: 608-613.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **PAI-2 (E-1): sc-166539**, our highly recommended monoclonal alternative to PAI-2 (H-70).