

PDE4A (H-166): sc-25810

BACKGROUND

Phosphodiesterases (PDE) hydrolyze cAMP to 5'AMP and thus play a critical role in the regulation of intracellular cAMP. Division of the PDE superfamily by sequence homology and enzymatic properties yields 11 PDE families. A unique upstream conserved region (UCR) characterizes the PDE4 family. Four separate genes (A-D) encode the PDE4 enzymes, and alternative splicing generates short or long isoforms of each gene. Long PDE4 isoforms contain both UCR1 and UCR2 while short PDE4 isoforms possess only UCR2. Both UCR domains are necessary for dimerization of PDE4 isoforms. The human PDE4A gene maps to chromosome 19p13.2 and spans 50 kilobases with 17 exons. The splice variants include isoforms PDE4A1-6.

REFERENCES

1. Bolger, G., et al. 1993. A family of human phosphodiesterases homologous to the dunce learning and memory gene product of *Drosophila melanogaster* are potential targets for antidepressant drugs. *Mol. Cell. Biol.* 13: 6558-6571.
2. Horton, Y.M., et al. 1995. Molecular cloning of a novel splice variant of human type IVA (PDE-IVA) cyclic AMP phosphodiesterase and localization of the gene to the p13.2-q12 region of human chromosome 19 [corrected]. *Biochem. J.* 308 (Pt 2):683-691.
3. SWISS-PROT/TrEMBL (P27815). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>.

CHROMOSOMAL LOCATION

Genetic locus: PDE4A (human) mapping to 19q13.2

SOURCE

PDE4A (H-166) is a rabbit polyclonal antibody raised against amino acids 721-886 mapping at the C-terminus of PDE4A of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PDE4A (H-166) is recommended for detection of PDE4A isoforms 1-5 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PDE4A siRNA (h): sc-41596, PDE4A shRNA Plasmid (h): sc-41596-SH and PDE4A shRNA (h) Lentiviral Particles: sc-41596-V.


Molecular Weight of PDE4A isoforms: 98/95/76/72/37 kDa.

Positive Controls: PDE4A (h): 293T Lysate: sc-113958, PDE4A (h4): 293T Lysate: sc-173505 or SH-SY5Y cell lysate: sc-3812.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/ 2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PDE4A (H-166): sc-25810. Western blot analysis of PDE4A expression in non-transfected: sc-117752 (A) and human PDE4A transfected: sc-173505 (B) 293T whole cell lysates.



PDE4A (H-166): sc-25810. Western blot analysis of PDE4A expression in non-transfected: sc-117752 (A) and human PDE4A transfected: sc-113958 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Dong, H., et al. 2010. Inhibition of PDE3, PDE4 and PDE7 potentiates glucocorticoid-induced apoptosis and overcomes glucocorticoid resistance in CEM T leukemic cells. *Biochem. Pharmacol.* 79: 321-329.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **PDE4A (H-7): sc-74428**, our highly recommended monoclonal alternative to PDE4A (H-166).