

melanotransferrin (H-70): sc-25832

BACKGROUND

Melanotransferrin is a member of the transferrin family of iron-binding proteins, which also includes serum transferrin, lactoferrin, and ovotransferrin, and it is highly expressed on melanoma cells. Melanotransferrin, also designated p97, shares a high degree of homology with transferrin, but does not play a significant role in the uptake of iron. Melanotransferrin utilizes a member of the low-density lipoprotein receptor family for transendothelial transport, which is not as efficient as the transport of transferrin through the corresponding transferrin receptor. The gene encoding human melanotransferrin maps to chromosome 3q29, and is predominantly expressed as either a membrane bound protein or a secreted form of the protein. Melanotransferrin is expressed in brain, where it may be involved in Alzheimer's disease. Melanotransferrin may also protect against membrane-lipid peroxidation, possess a metalloprotease activity, and possibly participate in intracellular adhesion. Further research will be necessary to fully elucidate the functions of this protein.

CHROMOSOMAL LOCATION

Genetic locus: MFI2 (human) mapping to 3q29; Mfi2 (mouse) mapping to 16 B2.

SOURCE

melanotransferrin (H-70) is a rabbit polyclonal antibody raised against amino acids 601-670 mapping near the C-terminus of melanotransferrin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

melanotransferrin (H-70) is recommended for detection of melanotransferrin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

melanotransferrin (H-70) is also recommended for detection of melanotransferrin in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for melanotransferrin siRNA (h): sc-41373, melanotransferrin siRNA (m): sc-41374, melanotransferrin shRNA Plasmid (h): sc-41373-SH, melanotransferrin shRNA Plasmid (m): sc-41374-SH, melanotransferrin shRNA (h) Lentiviral Particles: sc-41373-V and melanotransferrin shRNA (m) Lentiviral Particles: sc-41374-V.


Molecular Weight of melanotransferrin: 82 kDa.

Positive Controls: melanotransferrin (m): 293T Lysate: sc-121600, SK-MEL-28 cell lysate: sc-2236 or C32 whole cell lysate: sc-2205.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



melanotransferrin (H-70): sc-25832. Western blot analysis of melanotransferrin expression in non-transfected 293T: sc-117752 (A), mouse melanotransferrin transfected 293T: sc-121600 (B) and SK-MEL-28 (C) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **melanotransferrin (E-4): sc-271633** or **melanotransferrin (A-3): sc-271006**, our highly recommended monoclonal alternatives to melanotransferrin (H-70).