

# ADAM19 (A-16): sc-25985

## BACKGROUND

The ADAM (a disintegrin and metalloprotease) protein family, which includes over 30 membrane-anchored, glycosylated, Zn<sup>2+</sup> dependent proteases, plays a role in cell-cell and cell-matrix interface related processes, including fertilization, muscle fusion, secretion of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) and modulation of the neurogenic function of Notch and Delta. The ADAM proteins possess a signal-domain, a pro-domain, a metalloprotease domain, a disintegrin domain (integrin ligand), a cysteine-rich region, an epidermal growth factor-like domain, a transmembrane domain and a cytoplasmic tail. ADAMs are expressed in a wide range of mammalian tissues and several are abundantly expressed in the male reproductive tract. Expression of ADAM19, also designated Meltrin  $\beta$ , is highest in the peripheral nervous system during embryogenesis, but is also apparent in placenta, brain, heart, lung, leukocytes and SW480 cells. ADAM19 also serves as a dendritic cell marker. Truncation of ADAM19 in its cysteine-rich domain is necessary to exert its proteolytic activity on specific substrates, including  $\alpha_2$ -macroglobulin.

## REFERENCES

1. Wolfsberg, T.G., Primakoff, P., Myles, D.G. and White, J.M. 1995. ADAM, a novel family of membrane proteins containing a disintegrin and metalloprotease domain: multipotential functions in cell-cell and cell-matrix interactions. *J. Cell Biol.* 131: 275-278.
2. Stone, A.L., Kroeger, M. and Sang, Q.X. 1999. Structure-function analysis of the ADAM family of disintegrin-like and metalloproteinase-containing proteins. *J. Protein Chem.* 18: 447-465.
3. Fritsche, J., Moser, M., Faust, S., Peuker, A., Buttner, R., Andreesen, R. and Kreutz, M. 2000. Molecular cloning and characterization of a human metalloprotease disintegrin—a novel marker for dendritic cell differentiation. *Blood* 96: 732-739.
4. Primakoff, P. and Myles, D.G. 2000. The ADAM gene family: surface proteins with adhesion and protease activity. *Trends Genet.* 16: 83-87.
5. Wei, P., Zhao, Y.G., Zhuang, L., Ruben, S. and Sang, Q.X. 2001. Expression and enzymatic activity of human disintegrin and metalloproteinase ADAM19/meltrin  $\beta$ . *Biochem. Biophys. Res. Commun.* 280: 744-755.
6. Zhao, Y.G., Wei, P. and Sang, Q.X. 2001. Inhibitory antibodies against endopeptidase activity of human adamalysin 19. *Biochem. Biophys. Res. Commun.* 289: 288-294.

## CHROMOSOMAL LOCATION

Genetic locus: Adam19 (mouse) mapping to 11 B1.1.

## SOURCE

ADAM19 (A-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ADAM19 of mouse origin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-25985 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

ADAM19 (A-16) is recommended for detection of ADAM19 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ADAM19 (A-16) is also recommended for detection of ADAM19 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ADAM19 siRNA (m): sc-41418, ADAM19 shRNA Plasmid (m): sc-41418-SH and ADAM19 shRNA (m) Lentiviral Particles: sc-41418-V.

Molecular Weight of ADAM19 precursor: 115 kDa.

Molecular Weight of mature ADAM19: 87 kDa.

Positive Controls: IL-4 induced NIH/3T3 whole cell lysate or NIH/3T3 whole cell lysate: sc-2210.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **ADAM19 (14J12): sc-73687**, our highly recommended monoclonal alternative to ADAM19 (A-16).