

HuC (G-17): sc-26421

BACKGROUND

The Elav-like genes encode for a family of RNA-binding proteins. Elav, a *Drosophila* protein and the first described member, is expressed immediately after neuroblastic differentiation into neurons and is necessary for neuronal differentiation and maintenance. Several mammalian Elav-like proteins, designated HuC, HuD and Hel-N1, are also expressed in postmitotic neurons. An additional mammalian homolog, HuR, which is also designated HuA, is ubiquitously expressed and is also overexpressed in a wide variety of tumors. Characteristically, these homologs all contain three RNA recognition motifs (RRM), and they specifically bind to AU-rich elements (ARE) in the 3'-untranslated region of mRNAs transcripts. ARE sites target mRNA for rapid degradation and thereby regulate the expression levels of genes involved in cell growth and differentiation. When Elav-like proteins associate with these ARE sites this degradation is inhibited, leading to an increased stability of the corresponding transcript. Elav proteins function within the nucleus, and they are shuttled between the nucleus and cytoplasm by a nuclear export signal, which is a regulatory feature of the Elav-like proteins as it limits their accessibility to ARE sites.

CHROMOSOMAL LOCATION

Genetic locus: ELAVL3 (human) mapping to 19p13.2; Elavl3 (mouse) mapping to 9 A3.

SOURCE

HuC (G-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HuC of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26421 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HuC (G-17) is recommended for detection of HuC of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HuC (G-17) is also recommended for detection of HuC in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for HuC siRNA (h): sc-37833, HuC siRNA (m): sc-37834, HuC shRNA Plasmid (h): sc-37833-SH, HuC shRNA Plasmid (m): sc-37834-SH, HuC shRNA (h) Lentiviral Particles: sc-37833-V and HuC shRNA (m) Lentiviral Particles: sc-37834-V.


Molecular Weight of HuC: 38 kDa.

Positive Controls: SK-N-SH cell lysate: sc-2410, H4 cell lysate: sc-2408 or IMR-32 cell lysate: sc-2409.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



HuC (G-17): sc-26421. Western blot analysis of HuC expression in human brain tissue extract (A) and H4 (B), T98G (C) and IMR-32 (D) whole cell lysates.



HuC (G-17): sc-26421. Western blot analysis of HuC expression in SK-N-SH (A) and H4 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **HuC (D-6): sc-515624**, our highly recommended monoclonal alternative to HuC (G-17).