

GILZ (C-17): sc-26520

BACKGROUND

Glucocorticoid-induced leucine zipper (GILZ) is a leucine zipper protein expressed in normal lymphocytes from thymus, spleen and lymph nodes. GILZ is absent in nonlymphoid tissues including brain, liver and kidney. GILZ mediates the immunosuppressive effects of glucocorticoid hormones, and the expression of GILZ is induced in T cells by dexamethasone. GILZ protects T cells from an anti-CD3 antibody-induced apoptosis by inhibiting FAS and FAS ligand expression. GILZ interferes with Egr-2, Egr-3, NFAT/AP-1-inducible transcription factors and AP-1. The interaction of GILZ with c-Fos and c-Jun inhibits the binding of active AP-1 to its DNA consensus site *in vitro*. GILZ also binds NFκB subunits and inhibits the NFκB nuclear translocation. GILZ inhibits T cell receptor-induced interleukin-2/interleukin-2 receptor expression. The binding of GILZ to Raf-1 prevents Raf-MEK-ERK activation in the MAPK pathway. GILZ is expressed by normal macrophages in nonlymphoid tissues and by tumor-infiltrating macrophages in Burkitt lymphomas. The gene encoding human GILZ maps to chromosome Xq22.3.

REFERENCES

1. D'Adamio, F., Zollo, O., Moraca, R., Ayroldi, E., Bruscoli, S., Bartoli, A., Cannarile, L., Migliorati, G. and Riccardi, C. 1997. A new dexamethasone-induced gene of the leucine zipper family protects T lymphocytes from TCR/CD3-activated cell death. *Immunity* 7: 803-812.
2. Mittelstadt, P.R. and Ashwell, J.D. 2001. Inhibition of AP-1 by the glucocorticoid-inducible protein GILZ. *J. Biol. Chem.* 276: 29603-29610.
3. Ayroldi, E., Migliorati, G., Bruscoli, S., Marchetti, C., Zollo, O., Cannarile, L., D'Adamio, F. and Riccardi, C. 2001. Modulation of T cell activation by the glucocorticoid-induced leucine zipper factor via inhibition of nuclear factor κ B. *Blood* 98: 743-753.
4. Cannarile, L., Zollo, O., D'Adamio, F., Ayroldi, E., Marchetti, C., Tabilio, A., Bruscoli, S. and Riccardi, C. 2001. Cloning, chromosomal assignment and tissue distribution of human GILZ, a glucocorticoid hormone-induced gene. *Cell Death Differ.* 8: 201-203.
5. Ayroldi, E., Zollo, O., Macchiarulo, A., Di Marco, B., Marchetti, C. and Riccardi, C. 2002. Glucocorticoid-induced leucine zipper inhibits the Raf-extracellular signal-regulated kinase pathway by binding to Raf-1. *Mol. Cell. Biol.* 22: 7929-7941.
6. Berrebi, D., Bruscoli, S., Cohen, N., Foussat, A., Migliorati, G., Bouchet-Delbos, L., Maillot, M.C., Portier, A., Couderc, J., Galanaud, P., Peuchmaur, M., Riccardi, C. and Emilie, D. 2003. Synthesis of glucocorticoid-induced leucine zipper (GILZ) by macrophages: an anti-inflammatory and immunosuppressive mechanism shared by glucocorticoids and IL-10. *Blood* 101: 729-738.

CHROMOSOMAL LOCATION

Genetic locus: TSC22D3 (human) mapping to Xq22.3; Tsc22d3 (mouse) mapping to X F1.

SOURCE

GILZ (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of GILZ of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26520 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).


APPLICATIONS

GILZ (C-17) is recommended for detection of GILZ of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for GILZ siRNA (h): sc-43805, GILZ siRNA (m): sc-44809, GILZ shRNA Plasmid (h): sc-43805-SH, GILZ shRNA Plasmid (m): sc-44809-SH, GILZ shRNA (h) Lentiviral Particles: sc-43805-V and GILZ shRNA (m) Lentiviral Particles: sc-44809-V.

Molecular Weight of GILZ: 18 kDa.

DATA



GILZ (C-17): sc-26520. Western blot analysis of GILZ expression in non-transfected: sc-117752 (A) and human GILZ transfected: sc-111443 (B) 293T whole cell lysates



GILZ (C-17): sc-26520. Western blot analysis of GILZ expression in non-transfected: sc-117752 (A) and mouse GILZ transfected: sc-120485 (B) 293T whole cell lysates

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **GILZ (G-5): sc-133215** or **GILZ (D-2): sc-133216**, our highly recommended monoclonal alternative to GILZ (C-17).