

Rab 3B (V-13): sc-26555

BACKGROUND

The Ras-related superfamily of guanine nucleotide binding proteins, which includes the R-Ras, Rap, Ral/Rec and Rho/Rab subfamilies exhibit 30-60% homology with Ras p21. Accumulating data suggests an important role for Rab proteins, either in endocytosis or in biosynthetic protein transport. The transport of newly synthesized proteins from the endoplasmic reticulum to various stacks of the Golgi complex and to secretory vesicles involves at each stage the movement of carrier vesicles, a process that appears to involve Rab protein function. The possibility that Rab proteins might also direct the exocytosis from secretory vesicles to the plasma membrane is supported by the observation that in yeast, the SEC4 protein, which is 40% homologous to Rab proteins, is associated with secretory vesicles. At least eight members of the Rab subfamily have been identified, each of which is found at a particular stage of a membrane transport pathway.

REFERENCES

1. Zahraoui, A., et al. 1989. The human Rab genes encode a family of GTP-binding proteins related to yeast YPT1 and SEC4 products involved in secretion. *J. Biol. Chem.* 264: 12394-12401.
2. Chavrier, P., et al. 1992. The complexity of the Rab and Rho GTP-binding protein subfamilies revealed by a PCR cloning approach. *Gene* 112: 261-264.
3. Baldini, G., et al. 1992. Cloning of a Rab3 isotype predominately expressed in adipocytes. *Proc. Natl. Acad. Sci. USA* 89: 5049-5052.
4. Takizawa, P. and Malhotra, V. 1993. Coatomers and SNAREs in promoting membrane traffic. *Cell* 75: 593-596.
5. Novick, P. and Brennwald, P. 1993. Friends and family: the role of the Rab GTPases in vesicular traffic. *Cell* 75: 597-601.
6. Chen, Y., et al. 1993. Expression and localization of two low molecular weight GTP-binding proteins, Rab8 and Rab10, by epitope tag. *Proc. Natl. Acad. Sci. USA* 90: 6508-6512.
7. Torti, M., et al. 1993. Association of the low molecular weight GTP-binding protein rap2B with the cytoskeleton during platelet aggregation. *Proc. Natl. Acad. Sci. USA* 90: 7553-7557.
8. Karniguian, A., et al. 1993. Identification of small GTP-binding rab proteins in human platelets: Thrombin-induced phosphorylation of rab3B, rab6, and rab8 proteins. *Proc. Natl. Acad. Sci. USA* 90: 7647-7651.

CHROMOSOMAL LOCATIONS

Genetic locus: RAB3B (human) mapping to 1p32.3; Rab3b (mouse) mapping to 4 C7.

SOURCE

Rab 3B (V-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Rab 3B of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26555 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Rab 3B (V-13) is recommended for detection of Rab 3B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Rab 3B (V-13) is also recommended for detection of Rab 3B in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Rab 3B siRNA (h): sc-41814, Rab 3B siRNA (m): sc-41815, Rab 3B shRNA Plasmid (h): sc-41814-SH, Rab 3B shRNA Plasmid (m): sc-41815-SH, Rab 3B shRNA (h) Lentiviral Particles: sc-41814-V and Rab 3B shRNA (m) Lentiviral Particles: sc-41815-V.

Molecular Weight of Rab 3B: 25 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210 or IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Rab 3B (48-K2): sc-81911**, our highly recommended monoclonal alternative to Rab 3B (V-13).