

FCP1 (V-19): sc-26623

BACKGROUND

RNA polymerase II (RNAP II) comprises a 12 subunit protein complex that mediates transcription. RNAP II contains a unique carboxy terminal domain (CTD), which consists of 52 repeats of the consensus heptapeptide Tyr-Ser-Pro-Thr-Ser-Pro-Ser. The RNAP II isoform with an unphosphorylated CTD associates with transcription initiation complexes, whereas the isoform with a phosphorylated CTD is involved in transcription elongation. FCP1, also designated TFIIF-associated CTD phosphatase, is a protein phosphatase dedicated to the CTD of RNAP II that mediates the recycling of RNAP II during the transcription cycle. FCP1 specifically targets phosphorylated CTD Serine 2 and 5 with similar affinities. Phosphorylation of FCP1 itself is necessary for sufficient activity of the protein and its interaction with TFIIF. Upon activation, the α -helical carboxy terminus of FCP1 binds to RAP74 to form a complex. FCP1 also represses HIV-1 Tat-mediated transactivation and, therefore, may represent a specific target for modulation of Tat activity in infected cells.

REFERENCES

1. Licciardo, P., Ruggiero, L., Lania, L. and Majello, B. 2001. Transcription activation by targeted recruitment of the RNA polymerase II CTD phosphatase FCP1. *Nucleic Acids Res.* 29: 3539-3545.
2. Licciardo, P., Napolitano, G., Majello, B. and Lania, L. 2001. Inhibition of Tat transactivation by the RNA polymerase II CTD-phosphatase FCP1. *AIDS* 15: 301-307.
3. Mandal, S.S., Cho, H., Kim, S., Cabane, K. and Reinberg, D. 2002. FCP1, a phosphatase specific for the heptapeptide repeat of the largest subunit of RNA polymerase II, stimulates transcription elongation. *Mol. Cell. Biol.* 22: 7543-7552.
4. Lin, P.S., Dubois, M.F. and Dahmus, M.E. 2002. TFIIF-associating carboxyl-terminal domain phosphatase dephosphorylates phosphoserines 2 and 5 of RNA polymerase II. *J. Biol. Chem.* 277: 45949-45956.
5. Kamada, K., Roeder, R.G. and Burley, S.K. 2003. Molecular mechanism of recruitment of TFIIF- associating RNA polymerase C-terminal domain phosphatase (FCP1) by transcription factor IIF. *Proc. Natl. Acad. Sci. USA* 100: 2296-2299.
6. Friedl, E.M., Lane, W.S., Erdjument-Bromage, H., Tempst, P. and Reinberg, D. 2003. The C-terminal domain phosphatase and transcription elongation activities of FCP1 are regulated by phosphorylation. *Proc. Natl. Acad. Sci. USA* 100: 2328-2333.

CHROMOSOMAL LOCATION

Genetic locus: CTDPI (human) mapping to 18q23; Ctdp1 (mouse) mapping to 18 E3.

SOURCE

FCP1 (V-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FCP1 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26623 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

FCP1 (V-19) is recommended for detection of FCP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

FCP1 (V-19) is also recommended for detection of FCP1 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for FCP1 siRNA (h): sc-44888, FCP1 siRNA (m): sc-44889, FCP1 shRNA Plasmid (h): sc-44888-SH, FCP1 shRNA Plasmid (m): sc-44889-SH, FCP1 shRNA (h) Lentiviral Particles: sc-44888-V and FCP1 shRNA (m) Lentiviral Particles: sc-44889-V.

Molecular Weight of FCP1: 104 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **FCP1 (3G4): sc-293358**, our highly recommended monoclonal alternative to FCP1 (V-19).