

EMR1 (N-17): sc-26638

BACKGROUND

The epidermal growth factor (EGF)-TM7 family constitutes a group of class B G protein-coupled receptors, which includes CD97, EMR1 (EGF-like molecule containing mucin-like hormone receptor 1, designated F4/80 in mouse), EMR2, EMR3, FIRE, and ETL (1-3). These family members are characterized by an extended extracellular region with several N-terminal EGF domains, and are predominantly expressed on cells of the immune system (1-3). The EGF-TM7 protein family are encoded by a gene cluster on human chromosome 19p13 (1,3,4). The F4/80 molecule is solely expressed on the surface of macrophages and serves as a marker for mature macrophage tissues, including Kupffer cells in liver, splenic red pulp macrophages, brain microglia, gut lamina propria, and Langerhans cells in the skin. F4/80/EMR1 undergoes extensive N-linked glycosylation as well as some O-linked glycosylation (5,6). The function of F4/80/EMR1 is unclear, but it is speculated to be involved in macrophage adhesion events, cell migration, or as a G protein-coupled signaling component of macrophages.

REFERENCES

1. Baud, V., et al. 1995. EMR1, an unusual member in the family of hormone receptors with seven transmembrane segments. *Genomics* 26: 334-344.
2. Haidl, I.D., et al. 1996. The macrophage cell surface glycoprotein F4/80 is a highly glycosylated proteoglycan. *Eur. J. Immunol.* 26: 1139-1146.
3. Mander, T.H., et al. 1996. Development of microglia and macrophages in the postnatal rat pituitary. *Cell Tissue Res.* 286: 347-355.
4. Lin, H.H., et al. 2000. Human EMR2, a novel EGF-TM7 molecule on chromosome 19p13.1, is closely related to CD97. *Genomics* 67: 188-200.
5. Schaller, E., et al. 2002. Inactivation of the F4/80 glycoprotein in the mouse germ line. *Mol. Cell. Biol.* 22: 8035-8043.
6. Kwakkenbos, M.J., et al. 2002. The human EGF-TM7 family member EMR2 is a heterodimeric receptor expressed on myeloid cells. *J. Leukoc. Biol.* 71: 854-862.

CHROMOSOMAL LOCATION

Genetic locus: EMR1 (human) mapping to 19p13.3.

SOURCE

EMR1 (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of EMR1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26638 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

EMR1 (N-17) is recommended for detection of EMR1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for EMR1 siRNA (h): sc-72157, EMR1 shRNA Plasmid (h): sc-72157-SH and EMR1 shRNA (h) Lentiviral Particles: sc-72157-V.


Molecular Weight of EMR1: 160 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



EMR1 (N-17): sc-26638. Immunoperoxidase staining of formalin fixed, paraffin-embedded human fallopian tube tissue showing cytoplasmic and membrane staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **EMR1 (D-11): sc-365340**, our highly recommended monoclonal alternative to EMR1 (N-17).