

## SR-6 (P-20): sc-26669

### BACKGROUND

Serotonin (also designated 5-hydroxytryptamine or 5-HT) is a molecule that functions as a neurotransmitter, a hormone and a mitogen, and it is predominantly expressed in the gut, platelets and central nervous system (CNS). In the CNS, serotonin modulates several processes, including anxiety, sleep, appetite, behavior and drug abuse. In platelets and gut, serotonin plays a major role in cardiovascular function and motility of the gastrointestinal tract, respectively. Serotonin mediates its effects through several of G protein coupled receptors, designated 5-HT receptors or alternatively SR receptors. SR-3 is a ligand-gated ion channel, whereas all other known serotonin receptor subtypes are G protein-coupled receptors. Human SR-6 is one of the latest cloned receptors among the known SR receptors. SR-6 is expressed in the limbic region, which is involved in the control of mood and emotion and in nervous system diseases such as depression and Alzheimer's disease. The cellular mechanisms of SR-6 have not been clearly elucidated.

### REFERENCES

1. Maricq, A.V., et al. 1991. Primary structure and functional expression of the 5HT3 receptor, a serotonin-gated ion channel. *Science* 254: 432-437.
2. Kenakin, T.P., Bond, R.A., and Bonner, T.I. 1992. Definition of pharmacological receptors. *Pharmacol. Rev.* 44: 351-362.
3. Ruat, M., et al. 1993. Molecular cloning, characterization, and localization of a high-affinity serotonin receptor (5-HT7) activating cAMP formation. *Proc. Natl. Acad. Sci. USA* 90: 8547-8551.
4. Gelernter, J., et al. 1995. Assignment of the 5HT7 receptor gene (HTR7) to chromosome 10q and exclusion of genetic linkage with Tourette syndrome. *Genomics* 26: 207-209.
5. Weiss, B., et al. 1995. Assignment of a human homolog of the mouse Htr3 receptor gene to chromosome 11q23.1-q23.2. *Genomics* 29: 304-305.
6. Eglén, R.M., et al. 1995. Central 5-HT4 receptors. *Trends Pharmacol. Sci.* 16: 391-398.
7. Claeyens, S., et al. 1997. Assignment of 5-hydroxytryptamine receptor (HTR4) to human chromosome 5 bands q31→q33 by in situ hybridization. *Cytogenet. Cell Genet.* 78: 133-134.
8. Goppelt-Struebe, M., et al. 1998. Signaling pathways mediating induction of the early response genes prostaglandin G/H synthase-2 and egr-1 by serotonin via 5-HT2A receptors. *J. Cell Physiol.* 175: 341-347.

### CHROMOSOMAL LOCATIONS

Genetic locus: HTR6 (human) mapping to 1p36.13.

### SOURCE

SR-6 (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of SR-6 of human origin.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26669 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

SR-6 (P-20) is recommended for detection of SR-6 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SR-6 (P-20) is also recommended for detection of SR-6 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for SR-6 siRNA (h): sc-42244, SR-6 shRNA Plasmid (h): sc-42244-SH and SR-6 shRNA (h) Lentiviral Particles: sc-42244-V.

Molecular Weight of SR-6: 47 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.