



MMP-10 (P-16): sc-26697

BACKGROUND

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, Fibronectin, Laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-3, MMP-10 and MMP-11 (also designated stromelysin-1, -2 and -3) activate procollagenase. MMP-3 activation of procollagenase can occur via two pathways; direct activation by MMP-3 is slow while activation by MMP-3, in conjunction with tissue or plasma proteinases, is rapid. MMP-10 is expressed in small intestine, and at lower levels in lung and heart. MMP-11 is specifically expressed in stromal cells of breast carcinomas and contributes to epithelial cell malignancies.

REFERENCES

1. Saus, J., Quinones, S., Otani, Y., Nagase, H., Harris, E.D., Jr. and Kurkinen, M. 1988. The complete primary structure of human matrix metalloproteinase-3. Identity with stromelysin. *J. Biol. Chem.* 263: 6742-6745.
2. Suzuki, K., Enghild, J.J., Morodomi, T., Salvesen, G. and Nagase, H. 1990. Mechanisms of activation of tissue procollagenase by matrix metalloproteinase 3 (stromelysin). *Biochemistry* 29: 10261-10270.
3. Basset, P., Bellocq, J.P., Wolf, C., Stoll, I., Hutin, P., Limacher, J.M., Podhajcer, O.L., Chenard, M.P., Rio, M.C. and Chambon, P. 1990. A novel metalloproteinase gene specifically expressed in stromal cells of breast carcinomas. *Nature* 348: 699-704.
4. Birkedal-Hansen, H., Moore, W.G., Bodden, M.K., Windsor, L.J., Birkedal-Hansen, B., DeCarlo, A. and Engler, J.A. 1993. Matrix metalloproteinases: a review. *Crit. Rev. Oral Biol. Med.* 4: 197-250.
5. Reinemer, P., Grams, F., Huber, R., Kleine, T., Schnierer, S., Piper, M., Tschesche, H. and Bode, W. 1994. Structural implications for the role of the N-terminus in the "superactivation" of collagenases. A crystallographic study. *FEBS Lett.* 338: 227-233.
6. Knauper, V., Murphy, G. and Tschesche, H. 1996. Activation of human neutrophil procollagenase by stromelysin 2. *Eur. J. Biochem.* 235: 187-191.
7. Madlener, M. and Werner, S. 1997. cDNA cloning and expression of the gene encoding murine stromelysin-2 (MMP-10). *Gene* 202: 75-81.
8. Machein, U. and Conca, W. 1997. Expression of several matrix metalloproteinase genes in human monocytic cells. *Adv. Exp. Med. Biol.* 421: 247-251.

CHROMOSOMAL LOCATION

Genetic locus: *Mmp10* (mouse) mapping to 9 A1.

SOURCE

MMP-10 (P-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MMP-10 of mouse origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26697 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MMP-10 (P-16) is recommended for detection of MMP-10 of mouse and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MMP-10 siRNA (m): sc-41556, MMP-10 shRNA Plasmid (m): sc-41556-SH and MMP-10 shRNA (m) Lentiviral Particles: sc-41556-V.

Molecular Weight of MMP-10: 57 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.