

G_γ 3 (D-14): sc-26775

BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (e.g., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein α , β and γ polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their α subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Evidence, however, has established an important regulatory role for the $\beta\gamma$ subunits. It is becoming increasingly clear that different G protein complexes expressed in different tissues carry structurally distinct members of the γ as well as the α and β subunits and that preferential associations between members of subunit families increase G protein functional diversity.

CHROMOSOMAL LOCATION

Genetic locus: GNG3 (human) mapping to 11p12.3; Gng3 (mouse) mapping to 19 A.

SOURCE

G_γ 3 (D-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of G_γ 3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26775 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

G_γ 3 (D-14) is recommended for detection of G_γ 3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

G_γ 3 (D-14) is also recommended for detection of G_γ 3 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for G_γ 3 siRNA (h): sc-41776, G_γ 3 siRNA (m): sc-41777, G_γ 3 shRNA Plasmid (h): sc-41776-SH, G_γ 3 shRNA Plasmid (m): sc-41777-SH, G_γ 3 shRNA (h) Lentiviral Particles: sc-41776-V and G_γ 3 shRNA (m) Lentiviral Particles: sc-41777-V.


Molecular Weight of G_γ 3: 3-7 kDa.

Positive Controls: mouse cerebellum extract: sc-2403, rat brain extract: sc-2392 or mouse brain extract: sc-2253.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



G_γ 3 (D-14): sc-26775. Western blot analysis of G_γ 3 expression in mouse cerebellum (A) and rat brain (B) tissue extracts.



G_γ 3 (D-14): sc-26775. Western blot analysis of G_γ 3 expression in mouse brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try G_γ 3 (C-6): sc-393940, our highly recommended monoclonal alternative to G_γ 3 (D-14).