



Slit (dl-20): sc-26864

BACKGROUND

Drosophila melanogaster, a proven and effective model for studying developmental and cellular processes common to higher eukaryotes, contains a genome encoding approximately 13,600 genes, which were elucidated from more than 120 megabases of euchromatin. These genes are organized among chromosomes 2, 3, 4, X, and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes, which are categorized based on the type of protein for which they encode, represent six major classifications, including intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated), and other functional proteins. Morphogenesis and cell differentiation in *Drosophila* requires accurate control of cell division. The receptor Roundabout-1 (Robo1) and its ligand Slit influence axon guidance and central nervous system (CNS) patterning in both vertebrate and nonvertebrate systems. The Slit proteins exhibit a striking array of expression sites in non-neuronal tissues, including the urogenital system, limb primordia and developing eye.

REFERENCES

1. Lehner, C.F. 1991. Pulling the string: cell cycle regulation during *Drosophila* development. *Semin. Cell Biol.* 2: 223-31.
2. Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P.G., Scherer, S.E., Li, P.W., Hoskins, R.A., Galle, R.F., George, R.A., Lewis, S.E., Richards, S., Ashburner, M., Henderson, S.N., Sutton, G.G., Wortman, J.R., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
3. Mata, J., Curado, S., Ephrussi, A. and Rorth, P. 2000. Tribbles coordinates mitosis and morphogenesis in *Drosophila* by regulating string/CDC25 proteolysis. *Cell* 101: 511-22.
4. Connor, R.M. and Key, B. 2002. Expression and role of Roundabout-1 in embryonic *Xenopus* forebrain. *Dev Dyn.* 225: 22-34.
5. Piper, M. and Little, M. 2003. Movement through Slits: cellular migration via the Slit family. *Bioessays* 25: 32-38.
6. The Interactive Fly. <http://sdb.bio.purdue.edu/fly/aimain/1aahome.htm>.
<http://sdb.bio.purdue.edu/fly/aimain/6biochem.htm>

SOURCE

Slit (dl-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Slit of *Drosophila melanogaster* origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26864 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Slit (dl-20) is recommended for detection of Slit of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.