

# SMUG1 (N-19): sc-26878

## BACKGROUND

The base excision repair (BER) pathway removes incorrect bases (uracil) or damaged bases (3-methyladenine) from chromatin. Each BER enzyme system addresses a specific type of base damage. Uracil-DNA glycosylases, UNG2 and SMUG1 (single-strand selective monofunctional uracil DNA glycosylase) remove uracil from both double- and single-stranded DNA in nucleosomes (chromatin core particle). The uracil-excising enzyme family shares structural and functional conservation with minimal sequence conservation. The human SMUG1 gene maps to chromosome 12q13.13.

## REFERENCES

1. Haushalter, K.A., Todd Stukenberg, M.W., Kirschner, M.W. and Verdine, G.L. 1999. Identification of a new uracil-DNA glycosylase family by expression cloning using synthetic inhibitors. *Curr. Biol.* 9: 174-185.
2. Nilsen, H., Haushalter, K.A., Robins, P., Barnes, D.E., Verdine, G.L. and Lindahl, T. 2001. Excision of deaminated cytosine from the vertebrate genome: role of the SMUG1 uracil-DNA glycosylase. *EMBO J.* 20: 4278-4286.
3. Nilsen, H., Lindahl, T. and Verreault, A. 2002. DNA base excision repair of uracil residues in reconstituted nucleosome core particles. *EMBO J.* 21: 5943-5952.
4. Kavli, B., Sundheim, O., Akbari, M., Otterlei, M., Nilsen, H., Skorpen, F., Aas, P.A., Hagen, L., Krokan, H.E. and Slupphaug, G. 2002. hUNG2 is the major repair enzyme for removal of uracil from U:A matches, U:G mismatches, and U in single-stranded DNA, with hSMUG1 as a broad specificity backup. *J. Biol. Chem.* 277: 39926-39936.
5. LocusLink Report (LocusID: 2243). <http://www.ncbi.nlm.nih.gov/LocusLink/>

## CHROMOSOMAL LOCATION

Genetic locus: SMUG1 (human) mapping to 12q13.13; Smug1 (mouse) mapping to 15 F3.

## SOURCE

SMUG1 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SMUG1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26878 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

SMUG1 (N-19) is recommended for detection of SMUG1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SMUG1 (N-19) is also recommended for detection of SMUG1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SMUG1 siRNA (h): sc-106768, SMUG1 siRNA (m): sc-153643, SMUG1 shRNA Plasmid (h): sc-106768-SH, SMUG1 shRNA Plasmid (m): sc-153643-SH, SMUG1 shRNA (h) Lentiviral Particles: sc-106768-V and SMUG1 shRNA (m) Lentiviral Particles: sc-153643-V.


Molecular Weight of SMUG1: 34 kDa.

Positive Controls: SMUG1 (m): 293T Lysate: sc-123667.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



SMUG1 (N-19): sc-26878. Western blot analysis of SMUG1 expression in non-transfected: sc-117752 (A) and mouse SMUG1 transfected: sc-123667 (B) 293T whole cell lysates.



SMUG1 (N-19): sc-26878. Immunoperoxidase staining of formalin fixed, paraffin-embedded human ovary tissue showing nuclear and cytoplasmic staining of follicle cells and ovarian stroma cells.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.