# Dorsal (dF-19): sc-26908



The Power to Question

# **BACKGROUND**

Drosophila melanogaster is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. Drosophila genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. Morphogenesis and cell differentiation in Drosophila requires accurate control of cell division. The Drosophila Rel transcription factor Dorsal (DL) acts as the focal protein in the development of dorsoventral polarity in the developing fly. Dorsal activates and represses zygotic genes responsible for differentiation along the dorsoventral axis during the early stages of development. Dorsal and its inhibitor Cactus participate in a signal transduction pathway involved in several biologic processes, including embryonic pattern formation, immunity and muscle development.

# REFERENCES

- 1. Lehner, C.F. 1991. Pulling the string: cell cycle regulation during *Drosophila* development. Semin. Cell Biol. 2: 223-231.
- 2. Huang, A.M., Rusch, J., and Levine, M. 1997. An anteroposterior Dorsal gradient in the *Drosophila* embryo. Genes Dev. 11: 1963-1973.
- Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P., et al. 2000. The genome sequence of *Drosophila melanogaster*. Science. 287: 2185-2195.
- Mata, J., Curado, S., Ephrussi, A., and Rorth, P. 2000. Tribbles coordinates mitosis and morphogenesis in *Drosophila* by regulating string/Cdc25 proteolysis. Cell 101: 511-522.
- Bhaskar, V. and Courey, A.J. 2002. The MADF-BESS domain factor Dip3 potentiates synergistic activation by Dorsal and Twist. Gene 299: 173-184.
- Bolatto, C., Chifflet, S., Megighian, A., and Cantera, R. 2003. Synaptic activity modifies the levels of Dorsal and Cactus at the neuromuscular junction of *Drosophila*. J. Neurobiol. 54: 525-536.
- 7. The Interactive Fly. http://www.sdbonline.org/fly/aimain/1aahome.htm http://www.sdbonline.org/fly/aimain/6biochem.htm

# **SOURCE**

Dorsal (dF-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Dorsal of *Drosophila melanogaster* origin.

#### **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26908 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **APPLICATIONS**

Dorsal (dF-19) is recommended for detection of Dorsal of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

# **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com