



melanopsin (C-16): sc-26958

BACKGROUND

G protein coupled receptors (GPCRs) contain seven transmembrane helices and elicit G protein-mediated signaling cascades. The opsin family represents approximately 90 percent of all GPCRs and includes red, green, and blue-sensitive opsins, rhodopsin and melanopsin. Opsins consist of an apoprotein covalently linked to 11-cis-retinal that undergoes isomerization upon photon absorption. The photon-induced conformation change of opsin activates hundreds of G proteins. Mammalian melanopsin expression selectively occurs in the inner retina and not in the photoreceptor cells critical for vision. Melanopsin plays a nonessential role in the transduction of photic stimuli for light/dark entrainment. The gene encoding human melanopsin maps to chromosome 10q22.

REFERENCES

1. Fung, B.K., Hurley, J.B., and Stryer, L. 1980. Flow of information in the light-triggered cyclic nucleotide cascade of vision. *Proc. Natl. Acad. Sci. USA* 78: 152-156.
2. Hargrave, P.A., McDowell, J.H., Curtis, D.R., Wang, J.K., Juszczak, E., Fong, S.L., Rao, J.K., and Argos, P. 1983. The structure of bovine rhodopsin. *Biophys. Struct. Mech.* 9:235-244.
3. Iiri, T., Farfel, Z., and Bourne, H.R. 1998. G protein diseases furnish a model for the turn-on switch. *Nature* 394: 35-38.
4. Palczewski, K., Kumasaka, T., Hori, T., Behnke, C.A., Motoshima, H., Fox, B.A., Le Trong, I., Teller, D.C., Okada, T., Stenkamp, R.E., Yamamoto, M., and Miyano, M. 2000. Crystal structure of rhodopsin: a G protein-coupled receptor. *Science* 289: 739-745.
5. Provencio, I., Rodriguez, I.R., Jiang, G., Hayes, W.P., Moreira, E.F., and Rollag, M.D. 2000. A novel human opsin in the inner retina. *J. Neurosci.* 20:600-605.
6. Ruby, N.F., Brennan, T.J., Xie, X., Cao, V., Franken, P., Heller, H.C., and O'Hara, B.F. 2002. Role of melanopsin in circadian responses to light. *Science* 298: 2211-2213.

SOURCE

melanopsin (C-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of melanopsin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-26958 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

melanopsin (C-16) is recommended for detection of melanopsin of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for melanopsin siRNA (h): sc-40146.

Molecular Weight of melanopsin: 65 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.