

pyridoxal phosphatase (B-3): sc-271021

BACKGROUND

Pyridoxal phosphatase (PLPase) is an autoantigen comprising 296 amino acids. PLPase catalyzes the dephosphorylation of pyridoxal 5'-phosphate (the active form of vitamin B6) and exhibits a high level of expression various parts of the central nervous system, especially the brain. PLPase activity is catalyzed by haloacid dehalogenase (HAD), and it is the cofactor of both aromatic L-amino acid decarboxylase and glutamate decarboxylase. Autoantibodies against pyridoxal phosphatase show a strong correlation with certain types of cancer.

REFERENCES

- Choi, S.Y., et al. 1987. Brain pyridoxine-5-phosphate oxidase. Modulation of its by reaction with pyridoxal 5-phosphate and analogs. *J. Biol. Chem.* 262: 12013-12017.
- Jang, Y.M., et al. 2003. Human pyridoxal phosphatase. Molecular cloning, functional expression, and tissue distribution. *J. Biol. Chem.* 278: 50040-50046.
- Bøe, A.S., et al. 2004. Pyridoxal phosphatase system. *Br. J. Cancer* 91: 1508-1514.
- Kawai, S., et al. 2004. Cytosolic NADP phosphatases I and II from *Arthrobacter sp.* strain KM: implication in regulation of NAD⁺/NADP⁺ balance. *J. Basic Microbiol.* 44: 185-196.
- Kang, J.H., et al. 2004. Genomic organization, tissue distribution and deletion mutation of human pyridoxine 5'-phosphate oxidase. *Eur. J. Biochem.* 271: 2452-2461.
- Masse, P.G., et al. 2004. B-6 vitamers and 4-pyridoxic acid in the plasma, erythrocytes, and urine of postmenopausal women. *Am. J. Clin. Nutr.* 80: 946-951.
- Kim, D.W., et al. 2005. Inactivation of brain myo-inositol monophosphate phosphatase by pyridoxal-5'-phosphate. *J. Biochem. Mol. Biol.* 38: 58-64.
- Tirrell, I.M., et al. 2006. YZGD from *Paenibacillus thiaminolyticus*, a pyridoxal phosphatase of the HAD (haloacid dehalogenase) superfamily and a versatile member of the Nudix (nucleoside diphosphate x) hydrolase superfamily. *Biochem. J.* 394: 665-674.

CHROMOSOMAL LOCATION

Genetic locus: PDXP (human) mapping to 22q13.1.

SOURCE

pyridoxal phosphatase (B-3) is a mouse monoclonal antibody raised against amino acids 158-197 mapping within an internal region of pyridoxal phosphatase of human origin.

PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

pyridoxal phosphatase (B-3) is recommended for detection of pyridoxal phosphatase of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for pyridoxal phosphatase siRNA (h): sc-61425, pyridoxal phosphatase shRNA Plasmid (h): sc-61425-SH and pyridoxal phosphatase shRNA (h) Lentiviral Particles: sc-61425-V.


Molecular Weight of pyridoxal phosphatase: 32 kDa.

Positive Controls: pyridoxal phosphatase (h): 293T Lysate: sc-117114, Hep G2 cell lysate: sc-2227 or human brain extract: sc-364375.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



pyridoxal phosphatase (B-3): sc-271021. Western blot analysis of pyridoxal phosphatase expression in non-transfected: sc-117752 (A) and human pyridoxal phosphatase transfected: sc-117114 (B) 293T whole cell lysates.



pyridoxal phosphatase (B-3): sc-271021. Western blot analysis of pyridoxal phosphatase expression in human brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.