

OSM (T-18): sc-27104

BACKGROUND

Oncostatin M (OSM) is a glycoprotein that was originally isolated from the conditioned medium of U-937 human histiocytic leukemia cells that had been induced to differentiate into macrophage-like cells by treatment with phorbol 12-myristate 13-acetate. OSM inhibits the growth of a broad range of human tumor cell lines, but does not influence the growth of normal human fibroblasts. High-affinity binding sites for OSM have been detected on normal and tumor cells, and a receptor has been identified by chemical cross-linking studies. Expression of OSM is greatest in activated monocytic and lymphocytic cell lines and in normal adherent macrophages. Amino acid sequence analysis of OSM has revealed homology with leukemia inhibitory factor (LIF), granulocyte colony stimulating factor (G-CSF) and interleukin-6 (IL-6), all of which affect the growth and differentiation of a broad range of cell types, including those of hematopoietic origin.

REFERENCES

- Zarling, J.M., et al. 1986. Oncostatin M: a growth regulator produced by differentiated histiocytic lymphoma cells. *Proc. Natl. Acad. Sci. USA* 83: 9739-9743.
- Brown, T.J., et al. 1987. Purification and characterization of cytostatic lymphokines produced by activated human T lymphocytes. *J. Immunol.* 139: 2977-2983.
- Malik, N., et al. 1989. Molecular cloning, sequence analysis, and functional expression of a novel growth regulator, Oncostatin M. *Mol. Cell. Biol.* 9: 2847-2853.
- Linsley, P.S., et al. 1989. Identification and characterization of cellular receptors for the growth regulator, Oncostatin M. *J. Biol. Chem.* 264: 4282-4289.
- Horn, D., et al. 1990. Regulation of cell growth by recombinant Oncostatin M. *Growth Factors* 2: 157-165.

CHROMOSOMAL LOCATION

Genetic locus: OSM (human) mapping to 22q12.2; Osm (mouse) mapping to 11 A1.

SOURCE

OSM (T-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of OSM of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-27104 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

OSM (T-18) is recommended for detection of precursor and mature OSM of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for OSM siRNA (m): sc-39690, OSM shRNA Plasmid (m): sc-39690-SH and OSM shRNA (m) Lentiviral Particles: sc-39690-V.

Molecular Weight of OSM: 28 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



OSM (T-18): sc-27104. Western blot analysis of mouse recombinant OSM.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **OSM (A-9): sc-374039** or **OSM (H-2): sc-374040**, our highly recommended monoclonal alternatives to OSM (T-18).