# SANTA CRUZ BIOTECHNOLOGY, INC.

# EMAP II (G-2): sc-271115



# BACKGROUND

Endothelial monocyte-activating polypeptide (EMAP II), also known as small inducible cytokine subfamily E, member 1 (SCYE1), is a chemoattractant cytokine for monocytes and granulocytes that is inducible by apoptosis. TNF $\alpha$  treatment of murine meth A fibrosarcomas and B16 melanomas upregulates EMAP II mRNA production. The release of this cytokine renders the tumor-associated vasculature sensitive to tumor necrosis factor. EMAP II mRNA translates as a precursor protein, proEMAP II, which undergoes proteolysis to become the mature, biologically active cytokine. ProEMAP II may function in binding RNA as part of the tRNA synthetase complex in normal cells and in stimulating inflammatory responses after proteolytic cleavage in tumor cells.

# REFERENCES

- Knies, U.E., et al. 2000. Expression of EMAP II in the developing and adult mouse. Apoptosis 5: 141-151.
- Brabeck, C., et al. 2002. Expression of EMAP II by activated monocytes/ microglial cells in different regions of the rat hippocampus after trimethyltin-induced brain damage. Exp. Neurol. 177: 341-346.

## **CHROMOSOMAL LOCATION**

Genetic locus: AIMP1 (human) mapping to 4q24; Aimp1 (mouse) mapping to 3 G3.

# SOURCE

EMAP II (G-2) is a mouse monoclonal antibody raised against amino acids 221-312 mapping at the C-terminus of EMAP II of human origin.

# PRODUCT

Each vial contains 200  $\mu g\, lg G_1$  kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

# **APPLICATIONS**

EMAP II (G-2) is recommended for detection of precursor and mature EMAP II of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunohistochemistry (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for EMAP II siRNA (h): sc-61855, EMAP II siRNA (m): sc-61856, EMAP II shRNA Plasmid (h): sc-61855-SH, EMAP II shRNA Plasmid (m): sc-61856-SH, EMAP II shRNA (h) Lentiviral Particles: sc-61855-V and EMAP II shRNA (m) Lentiviral Particles: sc-61856-V.

Molecular Weight of EMAP II: 38-40 kDa.

Positive Controls: THP-1 cell lysate: sc-2238, WEHI-231 whole cell lysate: sc-2213 or A-10 cell lysate: sc-3806.

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG $\kappa$  BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA





FMAP II (G-2): sc-271115. Immunofluorescence staining

of methanol-fixed HeLa cells showing cytoplasmic

localization (A). Immunoperoxidase staining of formalin fixed. paraffin-embedded human brain tissue showing

cytoplasmic staining of neuronal and glial cells (B).

EMAP II (G-2): sc-271115. Western blot analysis of EMAP II expression in THP-1 ( $\mathbf{A}$ ), HT-1080 ( $\mathbf{B}$ ), WEHI-231 ( $\mathbf{C}$ ) and A-10 ( $\mathbf{D}$ ) whole cell lysates and rat testis tissue extract ( $\mathbf{E}$ ).

### **SELECT PRODUCT CITATIONS**

 Jang, S., et al. 2017. Suppression of AIMP1 protects cognition in Alzheimer's disease model mice 3xTg-AD. Neuroreport 28: 82-86.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.