

Endophilin B2 (C-10): sc-271129

BACKGROUND

The Endophilins comprise a family of proteins that associate with amphiphysin, synaptojanin and Dynamin and are implicated in presynaptic vesicle trafficking at nerve terminals. The expression patterns of the Endophilins are consistent with their cellular functions at the neuronal synapse. Endophilin B1 is a member of the B subgroup of the Endophilin family that is required for maintenance of mitochondrial morphology and for the regulation of the outer mitochondrial membrane dynamics. Endophilin B2 is also a member of the Endophilin B subgroup that is ubiquitously expressed but shows highest levels in brain, adult lung, ovary and spinal cord. Decreased levels of Endophilin B2 are found in Down syndrome and may reflect brain dysgenesis.

REFERENCES

- Pierrat, B., et al. 2001. SH3GLB, a new endophilin-related protein family featuring an SH3 domain. *Genomics* 71: 222-234.
- Modregger, J., et al. 2003. Characterization of Endophilin B1b, a brain-specific membrane-associated lysophosphatidic acid acyl transferase with properties distinct from Endophilin A1. *J. Biol. Chem.* 278: 4160-4167.
- Wang, M.Q., et al. 2003. Endophilins interact with Moloney murine leukemia virus $G_{\alpha\gamma}$ and modulate virion production. *J. Biol.* 3: 4.
- Engidawork, E., et al. 2003. Aberrant protein expression in cerebral cortex of fetus with Down syndrome. *Neuroscience* 122: 145-154.
- Karbowski, M., et al. 2004. Endophilin B1 is required for the maintenance of mitochondrial morphology. *J. Cell Biol.* 166: 1027-1039.
- Takahashi, Y., et al. 2005. Loss of Bif-1 suppresses Bax/Bak conformational change and mitochondrial apoptosis. *Mol. Cell. Biol.* 25: 9369-9382.
- Aramaki, Y., et al. 2005. Direct interaction between metastasis-associated protein 1 and Endophilin 3. *FEBS Lett.* 579: 3731-3736.

CHROMOSOMAL LOCATION

Genetic locus: SH3GLB2 (human) mapping to 9q34.11.

SOURCE

Endophilin B2 (C-10) is a mouse monoclonal antibody raised against amino acids 81-140 mapping near the N-terminus of Endophilin B2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Endophilin B2 (C-10) is recommended for detection of Endophilin B2 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Endophilin B2 siRNA (h): sc-60579, Endophilin B2 shRNA Plasmid (h): sc-60579-SH and Endophilin B2 shRNA (h) Lentiviral Particles: sc-60579-V.


Molecular Weight of Endophilin B2: 44 kDa.

Positive Controls: U-87 MG cell lysate: sc-2411 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Endophilin B2 (C-10): sc-271129. Western blot analysis of Endophilin B2 expression in U-87 MG (A) and HeLa (B) whole cell lysates.

Endophilin B2 (C-10): sc-271129. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human urinary bladder tissue showing cytoplasmic staining of urothelial cells (B).

RESEARCH USE

For research use only, not for use in diagnostic procedures.